

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
21 May 2004 (21.05.2004)

PCT

(10) International Publication Number  
**WO 2004/042389 A2**

(51) International Patent Classification<sup>7</sup>: **G01N 33/48**

RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,  
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:

PCT/EP2003/011877

(22) International Filing Date: 25 October 2003 (25.10.2003)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

02024993.4

8 November 2002 (08.11.2002) EP

(71) Applicant (for all designated States except US): **BAYER  
HEALTHCARE AG** [DE/DE]; 51368 Leverkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GOLZ, Stefan**  
[DE/DE]; Bückmannmühle 46, 45326 Essen (DE).  
**BRÜGGEMEIER, Ulf** [DE/DE]; Leysiefen 20, 42799  
Leichlingen (DE). **GEERTS, Andreas** [DE/DE]; Schuck-  
ertstr. 29, 42113 Wuppertal (DE).

(74) Common Representative: **BAYER HEALTHCARE**  
AG; Law and Patents, Patents and Licensing, 51368  
Leverkusen (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,  
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,  
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,  
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**

— as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii)) for the following designations AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH,  
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES,  
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH,  
PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN,  
TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO  
patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG,  
ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU,  
TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE,  
DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT,  
RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: **DIAGNOSTICS AND THERAPEUTICS FOR DISEASES ASSOCIATED WITH HUMAN PHOSPHODIESTERASE  
4D (PDE4D)**

(57) Abstract: The invention provides a human PDE4D which is associated with the disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders. The invention also provides assays for the identification of compounds useful in the treatment or prevention of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders. The invention also features compounds which bind to and/or activate or inhibit the activity of PDE4D as well as pharmaceutical compositions comprising such compounds.



**WO 2004/042389 A2**

Diagnostics and Therapeutics for Diseases Associated with Human  
Phosphodiesterase 4D (PDE4D)

5     Technical field of the invention

The present invention is in the field of molecular biology, more particularly, the present invention relates to nucleic acid sequences and amino acid sequences of a human PDE4D and its regulation for the treatment of disorders of the peripheral and  
10     central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in mammals.

Background of the invention

15     PDE4D is a member of the enzyme family of phosphodiesterases (PDEs) [Baecker PA et al., 1994], [Le Jeune IR et al., 2002], [Nemoz G et al., 1996], WO200160860, US5977305, WO200228999. PDEs catalyze the hydrolyzation of 3', 5' cyclic nucleotides. That results in the formation of the respective nucleoside 5' monophosphates. The cyclic nucleotides cAMP and cGMP serve as crucial second messengers in a  
20     number of cellular signaling pathways. The PDEs as well as the guanylyl and adenylyl cyclases, which synthesize the cyclic nucleotides, are important cellular components to regulate the concentration of cyclic nucleotides and, thus, to regulate the signal transduction pathways. Because of their central role in regulating second messenger levels PDEs have been considered chemotherapeutic targets and have  
25     been worked on extensively.

Several families of PDEs have been identified. The nomenclature system includes first a number that indicates the PDE family. To date, eleven families (PDE 1-11) are known which are classified by: (i) primary structure; (ii) substrate preference; (iii)  
30     response to different modulators; (iv) sensitivity to specific inhibitors; and (v) modes of regulation [Loughney and Ferguson, (1996)]. The number indicating the family is

- 2 -

followed by a capital letter, indicating a distinct gene, and the capital letter followed by a second number, indicating a specific splice variant or a specific transcript that utilizes a unique transcription initiation site.

5 PDEs show of the following structural features:

All mammalian PDEs identified to date possess a highly conserved region of 270-300 amino acids in the carboxy terminal half of the protein [Charbonneau, et al. (1986)]. Here, the catalytic site for cAMP and/or cGMP hydrolysis and two putative zinc binding sites as well as family specific determinants are located [Beavo, (1995);  
10 Francis, et al. (1994)]. The amino terminal regions of the various PDEs are highly variable and include other family specific determinants and diverse regulatory motifs such as: (i) calmodulin binding sites (PDE1); (ii) non-catalytic cyclic GMP binding sites (PDE2, PDE5, PDE6); (iii) membrane targeting sites (PDE4); (iv) hydrophobic membrane association sites (PDE3); and (v) phosphorylation sites for either the  
15 calmodulin-dependent kinase II (PDE1), the cAMP-dependent kinase (PDE1, PDE3, PDE4), or the cGMP dependent kinase (PDE5) [Beavo, (1995); Manganiello, et al. (1995); Conti, et al. (1995)].

Members of the PDE1 family are calcium-calmodulin dependent. The group is  
20 composed of at least three genes with several splicing variants [Kakkar, R. et al. (1999)]; PDE1A and PDE1B preferentially hydrolyze cGMP while PDE1C is dual-specific, it exhibits a high affinity for both cAMP and cGMP. In vitro experiments show regulation of some PDE1 species by phosphorylation, which decreases the affinity of the enzyme for calmodulin [Kakkar, (1999)]. PDE1s have been shown to  
25 be expressed in lung, heart and brain.

The PDE2 family is characterized as being specifically stimulated by cGMP [Loughney and Ferguson, supra]. PDE2 species have been found in cerebellum, neocortex, heart, kidney, lung, pulmonary artery, and skeletal muscle [Sadhu, K. et  
30 al. (1999)]. Only one gene has been identified, PDE2A. The respective PDE2A protein is specifically inhibited by erythro-9-(2-hydroxy-3-nonyl)adenine (EHNA).

Two genes have been identified in the PDE3 family, PDE3A and PDE3B, both having high affinity for both cAMP and cGMP, although the  $V_{\max}$  for cGMP hydrolysis is low enough that cGMP functions as a competitive inhibitor for cAMP hydrolysis. Enzymes in the PDE3 family are specifically inhibited by cGMP. PDE3 enzymes are specifically inhibited by milrinone and enoximone [Loughney and Ferguson, supra].

PDE4s are specific for cAMP hydrolysis. The family is comprised of four genes, PDE4A, PDE4B, PDE4C, and PDE4D. The genes give rise to multiple splice variants and are expressed in airway smooth muscle, the vascular endothelium, and all inflammatory cells. The enzymes can be activated by cAMP-dependent phosphorylation. Members of this family are specifically inhibited by the anti-depressant drug rolipram.

PDE5 is highly selective for cGMP [Turko, I.V. et al. (1998)]. Members of PDE5 family bind cGMP at non-catalytic sites [McAllister-Lucas, L.M. (1995)]. CGMP binding at non-catalytic sides has been suggested to be important for phosphorylation by cGMP-dependent protein kinase. PDE5 is highly expressed in vascular smooth muscle, platelets, lung, and kidney. Only one gene, PDE5A, has been identified.

PDE6s, the photoreceptor enzymes specifically hydrolyze cGMP [Loughney and Ferguson, supra]. PDE6s possess 2 regulatory high affinity cGMP binding sides. Genes include PDE6A and PDE6B (the protein products of which dimerize and bind two copies of a smaller  $\gamma$  inhibitory subunit to form rod PDE), in addition to PDE6C which associates with three smaller proteins to form cone PDE.

The PDE7 family effects cAMP hydrolysis but, in contrast to the PDE4 family, is not inhibited by rolipram [Loughney and Ferguson, supra]. Only one gene, PDE7A, has been identified. PDE7A gives rise to multiple splice variants. PDE7 mRNA can be found in several tissues but PDE7 protein expression appears to be restricted [Han, P.



- 4 -

et al. (1997); Perry, M.J. and G.A. Higgs (1998)]. Not much is known about the physiological function of PDE7.

5 The PDE8 family is closely related to the the PDE4 family. PDE8s have been shown to hydrolyze both cAMP and cGMP and are insensitive to inhibitors specific for PDEs 1-5. PDE8s are found in thyroid gland, testis, eye, liver, skeletal muscle, heart, kidney, ovary, and brain.

10 The PDE9 family preferentially hydrolyzes cAMP and is not sensitive to inhibition by rolipram, a PDE4-specific inhibitor, or isobutyl methyl xanthine (IBMX), a non-specific PDE inhibitor. PDE9 expression has been demonstrated in kidney, liver, lung, brain, spleen, and small intestine. Depending on nomenclature used, PDE9 is also referred to as PDE8, but is distinct from PDE8 mentioned above. To date, two genes have been identified in the PDE9 family.

15 PDE10 family members hydrolyze both cAMP and cGMP. PDE10s show expression in brain, thyroid and testis. [Soderling, S.H. et al. (1999); Fujishige, K. et al. (1999); Loughney, K. et al (1999)]

20 Members of the recently identified PDE11 family are also dualspecific. Interestingly, PDE11 splice variants exhibit different regulatory sequences in the N-terminal region. This suggests the possibility of differential regulation of PDE11s [Hetman JM, Robas N, Baxendale R, Fidock M, Phillips SC, Soderling SH, Beavo JA (2000)].

25 Increased PDE activity and decreased levels of cyclic nucleotides have been shown to be associated with many diseases. Furthermore, specific and non-specific inhibitors of several PDE protein families have been shown to be effective in treating such disorders. For example, the PDE4-specific inhibitor rolipram, mentioned above as an anti-depressant, inhibits lipopolysaccharide-induced expression of TNF- $\alpha$ , and  
30 has been effective in treating multiple sclerosis in an animal model. Other PDE4-specific inhibitors are being investigated for use as anti-inflammatory therapeutics,

- 5 -

and efficacy in attenuating the late asthmatic response to allergen challenge has been demonstrated [Harbinson, et al. (1997)]. Inhibitors specific for the PDE3 family have been approved for treatment of congestive heart failure. PDE5 inhibitors such as Sildenafil are in use for treatment of penile erectile dysfunction [Terrett, N. et al. (1996)]. PDE5-inhibitors are under investigation as agents for cardiovascular therapy [Perry, M.J. and G.A. Higgs (1998)].

PDEs cyclic nucleotide levels have been suggested to influence proliferation of different cell types [Conti et al. (1995)]. For example, growth of the prostatic carcinoma cell lines DU145 and LNCaP was inhibited by cAMP derivatives and PDE inhibitors [Bang, Y.J. et al. (1994)]. Furthermore, PDEs have been implemented to additional cancers.

Non-specific inhibitors, such as theophylline and pentoxifylline, are currently used in the treatment of respiratory and vascular disorders, respectively.

In summary, cAMP and cGMP play a central role in intracellular second messenger signaling. Furthermore, the value as pharmaceutical targets has been proven for several PDEs. Selective inhibitors have been developed as therapeutic agents for diseases such as cancer, heart failure, depression and sexual dysfunction. Thus, the identification of further disease implications of PDE species and their splice variants may lead to the development of specific inhibitors or modulators, or suggest new utilities for known compounds affecting PDEs. That in turn will provide additional pharmacological approaches to treat diseases and conditions in which alterations in cyclic nucleotide pathways are involved. This diseases may include, but are not limited to, infections such as bacterial, fungal, protozoan, and viral infections, particularly those caused by HIV viruses, cancers, allergies including asthma, cardiovascular diseases including acute heart failure, hypotension, hypertension, angina pectoris, myocardial infarction, hematological diseases, genito-urinary diseases including urinary incontinence and benign prostate hyperplasia, osteo-

porosis, and peripheral and central nervous system disorders including pain, Alzheimer's disease and Parkinson's disease.

#### **TaqMan-Technology / expression profiling**

5

TaqMan is a recently developed technique, in which the release of a fluorescent reporter dye from a hybridisation probe in real-time during a polymerase chain reaction (PCR) is proportional to the accumulation of the PCR product. Quantification is based on the early, linear part of the reaction, and by determining  
10 the threshold cycle (CT), at which fluorescence above background is first detected.

Gene expression technologies may be useful in several areas of drug discovery and development, such as target identification, lead optimization, and identification of mechanisms of action. The TaqMan technology can be used to compare differences  
15 between expression profiles of normal tissue and diseased tissue. Expression profiling has been used in identifying genes, which are up- or downregulated in a variety of diseases. An interesting application of expression profiling is temporal monitoring of changes in gene expression during disease progression and drug treatment or in patients versus healthy individuals. The premise in this approach is  
20 that changes in pattern of gene expression in response to physiological or environmental stimuli (e.g., drugs) may serve as indirect clues about disease-causing genes or drug targets. Moreover, the effects of drugs with established efficacy on global gene expression patterns may provide a guidepost, or a genetic signature, against which a new drug candidate can be compared.

25

#### **PDE4D**

The nucleotide sequence of PDE4D is accessible in public databases by the accession number Q08499 and is given in SEQ ID NO: 1. The amino acid sequence of PDE4D  
30 is depicted in SEQ ID NO: 2.

- 7 -

PDE4D is described as a phosphodiesterase [Baecker PA et al., 1994]. The phosphodiesterase PDE4D is published in [Le Jeune IR et al., 2002], WO 200160860, US 5977305 and WO 200228999. The expression of PDE4D in asthma relevant tissues and mononuclear cells was previously described [Nemoz G et al., 1996] [Le Jeune IR et al., 2002]. PDE4D shows the highest homology to other phosphodiesterases as shown in example 1.

### **Summary of the invention**

The invention relates to novel disease associations of PDE4D polypeptides and polynucleotides. The invention also relates to novel methods of screening for therapeutic agents for the treatment of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal. The invention also relates to pharmaceutical compositions for the treatment of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising a PDE4D polypeptide, a PDE4D polynucleotide, or regulators of PDE4D or modulators of PDE4D activity. The invention further comprises methods of diagnosing disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal.

### **Brief Description of the Drawings**

Fig. 1 shows the nucleotide sequence of a PDE4D polynucleotide (SEQ ID NO: 1).  
Fig. 2 shows the amino acid sequence of a PDE4D polypeptide (SEQ ID NO: 2).  
Fig. 3 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO: 3).  
Fig. 4 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO: 4).

- 8 -

Fig. 5 shows a nucleotide sequence useful as a probe to detect proteins of the invention (SEQ ID NO: 5).

### Detailed description of the invention

5

#### **Definition of terms**

An "oligonucleotide" is a stretch of nucleotide residues which has a sufficient number of bases to be used as an oligomer, amplicon or probe in a polymerase chain reaction (PCR). Oligonucleotides are prepared from genomic or cDNA sequence and are used to amplify, reveal, or confirm the presence of a similar DNA or RNA in a particular cell or tissue. Oligonucleotides or oligomers comprise portions of a DNA sequence having at least about 10 nucleotides and as many as about 35 nucleotides, preferably about 25 nucleotides.

15

"Probes" may be derived from naturally occurring or recombinant single- or double-stranded nucleic acids or may be chemically synthesized. They are useful in detecting the presence of identical or similar sequences. Such probes may be labeled with reporter molecules using nick translation, Klenow fill-in reaction, PCR or other methods well known in the art. Nucleic acid probes may be used in southern, northern or in situ hybridizations to determine whether DNA or RNA encoding a certain protein is present in a cell type, tissue, or organ.

20

A "fragment of a polynucleotide" is a nucleic acid that comprises all or any part of a given nucleotide molecule, the fragment having fewer nucleotides than about 6 kb, preferably fewer than about 1 kb.

25

"Reporter molecules" are radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents which associate with a particular nucleotide or amino acid sequence, thereby establishing the presence of a certain sequence, or allowing for the quantification of a certain sequence.

30

“Chimeric” molecules may be constructed by introducing all or part of the nucleotide sequence of this invention into a vector containing additional nucleic acid sequence which might be expected to change any one or several of the following PDE4D characteristics: cellular location, distribution, ligand-binding affinities, interchain  
5 affinities, degradation/turnover rate, signaling, etc.

“Active”, with respect to a PDE4D polypeptide, refers to those forms, fragments, or domains of a PDE4D polypeptide which retain the biological and/or antigenic  
10 activity of a PDE4D polypeptide.

“Naturally occurring PDE4D polypeptide” refers to a polypeptide produced by cells which have not been genetically engineered and specifically contemplates various polypeptides arising from post-translational modifications of the polypeptide including but not limited to acetylation, carboxylation, glycosylation, phos-  
15 phorylation, lipidation and acylation.

“Derivative” refers to polypeptides which have been chemically modified by techniques such as ubiquitination, labeling (see above), pegylation (derivatization with polyethylene glycol), and chemical insertion or substitution of amino acids such  
20 as ornithine which do not normally occur in human proteins.

“Conservative amino acid substitutions” result from replacing one amino acid with another having similar structural and/or chemical properties, such as the replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine  
25 with a serine.

“Insertions” or “deletions” are typically in the range of about 1 to 5 amino acids. The variation allowed may be experimentally determined by producing the peptide synthetically while systematically making insertions, deletions, or substitutions of  
30 nucleotides in the sequence using recombinant DNA techniques.

- 10 -

5 A "signal sequence" or "leader sequence" can be used, when desired, to direct the polypeptide through a membrane of a cell. Such a sequence may be naturally present on the polypeptides of the present invention or provided from heterologous sources by recombinant DNA techniques.

10 An "oligopeptide" is a short stretch of amino acid residues and may be expressed from an oligonucleotide. Oligopeptides comprise a stretch of amino acid residues of at least 3, 5, 10 amino acids and at most 10, 15, 25 amino acids, typically of at least 9 to 13 amino acids, and of sufficient length to display biological and/or antigenic activity.

15 "Inhibitor" is any substance which retards or prevents a chemical or physiological reaction or response. Common inhibitors include but are not limited to antisense molecules, antibodies, and antagonists.

"Standard expression" is a quantitative or qualitative measurement for comparison. It is based on a statistically appropriate number of normal samples and is created to use as a basis of comparison when performing diagnostic assays, running clinical trials, or following patient treatment profiles.

25 "Animal" as used herein may be defined to include human, domestic (e.g., cats, dogs, etc.), agricultural (e.g., cows, horses, sheep, etc.) or test species (e.g., mouse, rat, rabbit, etc.).

A "PDE4D polynucleotide", within the meaning of the invention, shall be understood as being a nucleic acid molecule selected from a group consisting of

30 (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2,

- 11 -

- (ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1, ...
- (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1,
- 5 (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii); and
- (v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code;

10

wherein the polypeptide encoded by said nucleic acid molecule has PDE4D activity.

A "PDE4D polypeptide", within the meaning of the invention, shall be understood as being a polypeptide selected from a group consisting of

15

- (i) polypeptides having the sequence of SEQ ID NO: 2,
- (ii) polypeptides comprising the sequence of SEQ ID NO: 2,
- 20 (iii) polypeptides encoded by PDE4D polynucleotides; and
- (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% homology with a polypeptide of (i), (ii), or (iii);

25

wherein said polypeptide has PDE4D activity.

The nucleotide sequences encoding a PDE4D (or their complement) have numerous applications in techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use in the construction of oligomers for PCR, use for chromosome and gene mapping, use in the recombinant  
30 production of PDE4D, and use in generation of antisense DNA or RNA, their



- 12 -

chemical analogs and the like. Uses of nucleotides encoding a PDE4D disclosed herein are exemplary of known techniques and are not intended to limit their use in any technique known to a person of ordinary skill in the art. Furthermore, the nucleotide sequences disclosed herein may be used in molecular biology techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, e.g., the triplet genetic code, specific base pair interactions, etc.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of PDE4D - encoding nucleotide sequences may be produced. Some of these will only bear minimal homology to the nucleotide sequence of the known and naturally occurring PDE4D. The invention has specifically contemplated each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring PDE4D, and all such variations are to be considered as being specifically disclosed.

Although the nucleotide sequences which encode a PDE4D, its derivatives or its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring PDE4D polynucleotide under stringent conditions, it may be advantageous to produce nucleotide sequences encoding PDE4D polypeptides or its derivatives possessing a substantially different codon usage. Codons can be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic expression host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding a PDE4D polypeptide and/or its derivatives without altering the encoded amino acid sequence include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

Nucleotide sequences encoding a PDE4D polypeptide may be joined to a variety of other nucleotide sequences by means of well established recombinant DNA techniques. Useful nucleotide sequences for joining to PDE4D polynucleotides include an assortment of cloning vectors such as plasmids, cosmids, lambda phage derivatives, phagemids, and the like. Vectors of interest include expression vectors, replication vectors, probe generation vectors, sequencing vectors, etc. In general, vectors of interest may contain an origin of replication functional in at least one organism, convenient restriction endonuclease sensitive sites, and selectable markers for one or more host cell systems.

Another aspect of the subject invention is to provide for PDE4D-specific hybridization probes capable of hybridizing with naturally occurring nucleotide sequences encoding PDE4D. Such probes may also be used for the detection of similar PDE encoding sequences and should preferably show at least 40% nucleotide identity to PDE4D polynucleotides. The hybridization probes of the subject invention may be derived from the nucleotide sequence presented as SEQ ID NO: 1 or from genomic sequences including promoter, enhancers or introns of the native gene. Hybridization probes may be labelled by a variety of reporter molecules using techniques well known in the art.

It will be recognized that many deletional or mutational analogs of PDE4D polynucleotides will be effective hybridization probes for PDE4D polynucleotides. Accordingly, the invention relates to nucleic acid sequences that hybridize with such PDE4D encoding nucleic acid sequences under stringent conditions.

"Stringent conditions" refers to conditions that allow for the hybridization of substantially related nucleic acid sequences. For instance, such conditions will generally allow hybridization of sequence with at least about 85% sequence identity, preferably with at least about 90% sequence identity, more preferably with at least about 95% sequence identity. Hybridization conditions and probes can be adjusted in well-characterized ways to achieve selective hybridization of human-derived

- 14 -

probes. Stringent conditions, within the meaning of the invention are 65°C in a buffer containing 1 mM EDTA, 0.5 M NaHPO<sub>4</sub> (pH 7.2), 7 % (w/v) SDS.

5 Nucleic acid molecules that will hybridize to PDE4D polynucleotides under stringent conditions can be identified functionally. Without limitation, examples of the uses for hybridization probes include: histochemical uses such as identifying tissues that express PDE4D; measuring mRNA levels, for instance to identify a sample's tissue type or to identify cells that express abnormal levels of PDE4D; and detecting polymorphisms of PDE4D.

10 PCR provides additional uses for oligonucleotides based upon the nucleotide sequence which encodes PDE4D. Such probes used in PCR may be of recombinant origin, chemically synthesized, or a mixture of both. Oligomers may comprise discrete nucleotide sequences employed under optimized conditions for identification of PDE4D in specific tissues or diagnostic use. The same two oligomers, a nested set of oligomers, or even a degenerate pool of oligomers may be employed under less  
15 stringent conditions for identification of closely related DNAs or RNAs.

Rules for designing polymerase chain reaction (PCR) primers are now established, as  
20 reviewed by PCR Protocols. Degenerate primers, i.e., preparations of primers that are heterogeneous at given sequence locations, can be designed to amplify nucleic acid sequences that are highly homologous to, but not identical with PDE4D. Strategies are now available that allow for only one of the primers to be required to specifically hybridize with a known sequence. For example, appropriate nucleic acid  
25 primers can be ligated to the nucleic acid sought to be amplified to provide the hybridization partner for one of the primers. In this way, only one of the primers need be based on the sequence of the nucleic acid sought to be amplified.

30 PCR methods for amplifying nucleic acid will utilize at least two primers. One of these primers will be capable of hybridizing to a first strand of the nucleic acid to be amplified and of priming enzyme-driven nucleic acid synthesis in a first direction.

The other will be capable of hybridizing the reciprocal sequence of the first strand (if the sequence to be amplified is single stranded, this sequence will initially be hypothetical, but will be synthesized in the first amplification cycle) and of priming nucleic acid synthesis from that strand in the direction opposite the first direction and  
5 towards the site of hybridization for the first primer. Conditions for conducting such amplifications, particularly under preferred stringent hybridization conditions, are well known.

Other means of producing specific hybridization probes for PDE4D include the  
10 cloning of nucleic acid sequences encoding PDE4D or PDE4D derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate reporter molecules.

15 It is possible to produce a DNA sequence, or portions thereof, entirely by synthetic chemistry. After synthesis, the nucleic acid sequence can be inserted into any of the many available DNA vectors and their respective host cells using techniques which are well known in the art. Moreover, synthetic chemistry may be used to introduce  
20 mutations into the nucleotide sequence. Alternately, a portion of sequence in which a mutation is desired can be synthesized and recombined with longer portion of an existing genomic or recombinant sequence.

PDE4D polynucleotides may be used to produce a purified oligo- or polypeptide  
25 using well known methods of recombinant DNA technology. The oligopeptide may be expressed in a variety of host cells, either prokaryotic or eukaryotic. Host cells may be from the same species from which the nucleotide sequence was derived or from a different species. Advantages of producing an oligonucleotide by recombinant DNA technology include obtaining adequate amounts of the protein for  
30 purification and the availability of simplified purification procedures.

### Quantitative determinations of nucleic acids

An important step in the molecular genetic analysis of human disease is often the enumeration of the copy number of a nucleic acid or the relative expression of a gene in particular tissues.

Several different approaches are currently available to make quantitative determinations of nucleic acids. Chromosome-based techniques, such as comparative genomic hybridization (CGH) and fluorescent in situ hybridization (FISH) facilitate efforts to cytogenetically localize genomic regions that are altered in tumor cells. Regions of genomic alteration can be narrowed further using loss of heterozygosity analysis (LOH), in which disease DNA is analyzed and compared with normal DNA for the loss of a heterozygous polymorphic marker. The first experiments used restriction fragment length polymorphisms (RFLPs) [Johnson, (1989)], or hyper-variable minisatellite DNA [Barnes, 2000]. In recent years LOH has been performed primarily using PCR amplification of microsatellite markers and electrophoresis of the radio labelled [Jeffreys, (1985)] or fluorescently labelled PCR products [Weber, (1990)] and compared between paired normal and disease DNAs.

A number of other methods have also been developed to quantify nucleic acids [Gergen, (1992)]. More recently, PCR and RT-PCR methods have been developed which are capable of measuring the amount of a nucleic acid in a sample. One approach, for example, measures PCR product quantity in the log phase of the reaction before the formation of reaction products plateaus [Thomas, (1980)].

A gene sequence contained in all samples at relatively constant quantity is typically utilized for sample amplification efficiency normalization. This approach, however, suffers from several drawbacks. The method requires that each sample has equal input amounts of the nucleic acid and that the amplification efficiency between samples is identical until the time of analysis. Furthermore, it is difficult using the conventional methods of PCR quantitation such as gel electrophoresis or plate

- 17 -

capture hybridization to determine that all samples are in fact analyzed during the log phase of the reaction as required by the method.

5 Another method called quantitative competitive (QC)-PCR, as the name implies, relies on the inclusion of an internal control competitor in each reaction [Piatak, (1993), BioTechniques]. The efficiency of each reaction is normalized to the internal competitor. A known amount of internal competitor is typically added to each sample. The unknown target PCR product is compared with the known competitor PCR product to obtain relative quantitation. A difficulty with this general approach  
10 lies in developing an internal control that amplifies with the same efficiency than the target molecule.

#### *5' Fluorogenic Nuclease Assays*

15 Fluorogenic nuclease assays are a real time quantitation method that uses a probe to monitor formation of amplification product. The basis for this method of monitoring the formation of amplification product is to measure continuously PCR product accumulation using a dual-labelled fluorogenic oligonucleotide probe, an approach frequently referred to in the literature simply as the "TaqMan method" [Piatak,  
20 (1993), Science; Heid, (1996); Gibson, (1996); Holland. (1991)].

The probe used in such assays is typically a short (about 20-25 bases) oligonucleotide that is labeled with two different fluorescent dyes. The 5' terminus of the probe is attached to a reporter dye and the 3' terminus is attached to a quenching dye,  
25 although the dyes could be attached at other locations on the probe as well. The probe is designed to have at least substantial sequence complementarity with the probe binding site. Upstream and downstream PCR primers which bind to flanking regions of the locus are added to the reaction mixture. When the probe is intact, energy transfer between the two fluorophors occurs and the quencher quenches  
30 emission from the reporter. During the extension phase of PCR, the probe is cleaved by the 5' nuclease activity of a nucleic acid polymerase such as Taq polymerase,

- 18 -

thereby releasing the reporter from the oligonucleotide-quencher and resulting in an increase of reporter emission intensity which can be measured by an appropriate detector.

5 One detector which is specifically adapted for measuring fluorescence emissions such as those created during a fluorogenic assay is the ABI 7700 or 4700 HT manufactured by Applied Biosystems, Inc. in Foster City, Calif. The ABI 7700 uses fiber optics connected with each well in a 96-or 384 well PCR tube arrangement. The instrument includes a laser for exciting the labels and is capable of measuring  
10 the fluorescence spectra intensity from each tube with continuous monitoring during PCR amplification. Each tube is re-examined every 8.5 seconds.

Computer software provided with the instrument is capable of recording the fluorescence intensity of reporter and quencher over the course of the amplification.  
15 The recorded values will then be used to calculate the increase in normalized reporter emission intensity on a continuous basis. The increase in emission intensity is plotted versus time, i.e., the number of amplification cycles, to produce a continuous measure of amplification. To quantify the locus in each amplification reaction, the amplification plot is examined at a point during the log phase of product  
20 accumulation. This is accomplished by assigning a fluorescence threshold intensity above background and determining the point at which each amplification plot crosses the threshold (defined as the threshold cycle number or Ct). Differences in threshold cycle number are used to quantify the relative amount of PCR target contained within each tube. Assuming that each reaction functions at 100% PCR efficiency, a  
25 difference of one Ct represents a two-fold difference in the amount of starting template. The fluorescence value can be used in conjunction with a standard curve to determine the amount of amplification product present.

*Non-Probe-Based Detection Methods*

A variety of options are available for measuring the amplification products as they are formed. One method utilizes labels, such as dyes, which only bind to double stranded DNA. In this type of approach, amplification product (which is double  
5 stranded) binds dye molecules in solution to form a complex. With the appropriate dyes, it is possible to distinguish between dye molecules free in solution and dye molecules bound to amplification product. For example, certain dyes fluoresce only when bound to amplification product. Examples of dyes which can be used in  
10 methods of this general type include, but are not limited to, Syber Green.TM. and Pico Green from Molecular Probes, Inc. of Eugene, Oreg., ethidium bromide, propidium iodide, chromomycin, acridine orange, Hoechst 33258, Toto-1, Yoyo-1, DAPI (4',6-diamidino-2-phenylindole hydrochloride).

15 Another real time detection technique measures alteration in energy fluorescence energy transfer between fluorophors conjugated with PCR primers [Livak, (1995)].

*Probe-Based Detection Methods*

20 These detection methods involve some alteration to the structure or conformation of a probe hybridized to the locus between the amplification primer pair. In some instances, the alteration is caused by the template-dependent extension catalyzed by a nucleic acid polymerase during the amplification process. The alteration generates a detectable signal which is an indirect measure of the amount of amplification product  
25 formed.

For example, some methods involve the degradation or digestion of the probe during the extension reaction. These methods are a consequence of the 5'-3' nuclease activity associated with some nucleic acid polymerases. Polymerases having this  
30 activity cleave mononucleotides or small oligonucleotides from an oligonucleotide probe annealed to its complementary sequence located within the locus.



- 20 -

The 3' end of the upstream primer provides the initial binding site for the nucleic acid polymerase. As the polymerase catalyzes extension of the upstream primer and encounters the bound probe, the nucleic acid polymerase displaces a portion of the 5' end of the probe and through its nuclease activity cleaves mononucleotides or oligonucleotides from the probe.

The upstream primer and the probe can be designed such that they anneal to the complementary strand in close proximity to one another. In fact, the 3' end of the upstream primer and the 5' end of the probe may abut one another. In this situation, extension of the upstream primer is not necessary in order for the nucleic acid polymerase to begin cleaving the probe. In the case in which intervening nucleotides separate the upstream primer and the probe, extension of the primer is necessary before the nucleic acid polymerase encounters the 5' end of the probe. Once contact occurs and polymerization continues, the 5'-3' exonuclease activity of the nucleic acid polymerase begins cleaving mononucleotides or oligonucleotides from the 5' end of the probe. Digestion of the probe continues until the remaining portion of the probe dissociates from the complementary strand.

In solution, the two end sections can hybridize with each other to form a hairpin loop. In this conformation, the reporter and quencher dye are in sufficiently close proximity that fluorescence from the reporter dye is effectively quenched by the quencher dye. Hybridized probe, in contrast, results in a linearized conformation in which the extent of quenching is decreased. Thus, by monitoring emission changes for the two dyes, it is possible to indirectly monitor the formation of amplification product.

### *Probes*

The labeled probe is selected so that its sequence is substantially complementary to a segment of the test locus or a reference locus. As indicated above, the nucleic acid

site to which the probe binds should be located between the primer binding sites for the upstream and downstream amplification primers.

### *Primers*

5

The primers used in the amplification are selected so as to be capable of hybridizing to sequences at flanking regions of the locus being amplified. The primers are chosen to have at least substantial complementarity with the different strands of the nucleic acid being amplified. When a probe is utilized to detect the formation of amplification products, the primers are selected in such that they flank the probe, i.e. are located upstream and downstream of the probe.

10

The primer must have sufficient length so that it is capable of priming the synthesis of extension products in the presence of an agent for polymerization. The length and composition of the primer depends on many parameters, including, for example, the temperature at which the annealing reaction is conducted, proximity of the probe binding site to that of the primer, relative concentrations of the primer and probe and the particular nucleic acid composition of the probe. Typically the primer includes 15-30 nucleotides. However, the length of the primer may be more or less depending on the complexity of the primer binding site and the factors listed above.

15

20

### *Labels for Probes and Primers*

The labels used for labeling the probes or primers of the current invention and which can provide the signal corresponding to the quantity of amplification product can take a variety of forms. As indicated above with regard to the 5' fluorogenic nuclease method, a fluorescent signal is one signal which can be measured. However, measurements may also be made, for example, by monitoring radioactivity, colorimetry, absorption, magnetic parameters, or enzymatic activity. Thus, labels which can be employed include, but are not limited to, fluorophores, chromophores,

25

30

- 22 -

radioactive isotopes, electron dense reagents, enzymes, and ligands having specific binding partners (e.g., biotin-avidin).

Monitoring changes in fluorescence is a particularly useful way to monitor the accumulation of amplification products. A number of labels useful for attachment to probes or primers are commercially available including fluorescein and various fluorescein derivatives such as FAM, HEX, TET and JOE (all which are available from Applied Biosystems, Foster City, Calif.); lucifer yellow, and coumarin derivatives.

Labels may be attached to the probe or primer using a variety of techniques and can be attached at the 5' end, and/or the 3' end and/or at an internal nucleotide. The label can also be attached to spacer arms of various sizes which are attached to the probe or primer. These spacer arms are useful for obtaining a desired distance between multiple labels attached to the probe or primer.

In some instances, a single label may be utilized; whereas, in other instances, such as with the 5' fluorogenic nuclease assays for example, two or more labels are attached to the probe. In cases wherein the probe includes multiple labels, it is generally advisable to maintain spacing between the labels which is sufficient to permit separation of the labels during digestion of the probe through the 5'-3' nuclease activity of the nucleic acid polymerase.

#### **Patients Exhibiting Symptoms of Disease**

A number of diseases are associated with changes in the copy number of a certain gene. For patients having symptoms of a disease, the real-time PCR method can be used to determine if the patient has copy number alterations which are known to be linked with diseases that are associated with the symptoms the patient has.

## PDE4D expression

### *PDE4D fusion proteins*

5 Fusion proteins are useful for generating antibodies against PDE4D polypeptides and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with portions of PDE4D polypeptides. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such  
10 methods are well known in the art and also can be used as drug screens.

A PDE4D fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment can comprise at least 54, 75, 100, 125, 139, 150, 175, 200, 225, 250, 275, 300, 325 or 350 contiguous amino acids  
15 of SEQ ID NO: 2 or of a biologically active variant, such as those described above. The first polypeptide segment also can comprise full-length PDE4D.

The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include, but are not limited to  
20  $\beta$ -galactosidase,  $\beta$ -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including  
25 histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions. A fusion protein also can be engineered to contain a cleavage site located adjacent to the PDE4D.

*Preparation of Polynucleotides*

A naturally occurring PDE4D polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides  
5 can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique for obtaining a polynucleotide can be used to obtain isolated PDE4D polynucleotides.  
10 For example, restriction enzymes and probes can be used to isolate polynucleotide fragments which comprise PDE4D nucleotide sequences. Isolated polynucleotides are in preparations which are free or at least 70, 80, or 90% free of other molecules.

PDE4D cDNA molecules can be made with standard molecular biology techniques,  
15 using PDE4D mRNA as a template. PDE4D cDNA molecules can thereafter be replicated using molecular biology techniques known in the art. An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or cDNA as a template.

20 Alternatively, synthetic chemistry techniques can be used to synthesize PDE4D polynucleotides. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode PDE4D having, for example, an amino acid sequence shown in SEQ ID NO: 2 or a biologically active variant thereof.

*Extending Polynucleotides*

Various PCR-based methods can be used to extend nucleic acid sequences encoding human PDE4D, for example to detect upstream sequences of PDE4D gene such as promoters and regulatory elements. For example, restriction-site PCR uses universal  
30 primers to retrieve unknown sequence adjacent to a known locus. Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific

- 25 -

to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

5

Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region. Primers can be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72°C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

10

Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. In this method, multiple restriction enzyme digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

15

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries can be useful for extension of sequence into 5' non-transcribed regulatory regions.

20

Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser

25

30

- 26 -

activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate equipment and software (e.g., GENOTYPER and Sequence NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and  
5 electronic data display can be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

#### *Obtaining Polypeptides*

10

PDE4D can be obtained, for example, by purification from human cells, by expression of PDE4D polynucleotides, or by direct chemical synthesis.

#### *Protein Purification*

15

PDE4D can be purified from any human cell which expresses the enzyme, including those which have been transfected with expression constructs which express PDE4D. A purified PDE4D is separated from other compounds which normally associate with PDE4D in the cell, such as certain proteins, carbohydrates, or lipids, using methods  
20 well-known in the art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis.

#### *Expression of PDE4D Polynucleotides*

25

To express PDE4D, PDE4D polynucleotides can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art can be used to construct expression vectors containing sequences encoding  
30 PDE4D and appropriate transcriptional and translational control elements. These

- 27 -

methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination.

5 A variety of expression vector/host systems can be utilized to contain and express sequences encoding PDE4D. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (*e.g.*, baculovirus), plant cell systems transformed with virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmids), or animal cell systems.

15 The control elements or regulatory sequences are those non-translated regions of the vector - enhancers, promoters, 5' and 3' untranslated regions -- which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (*e.g.*, heat shock, RUBISCO, and storage protein genes) or from plant viruses (*e.g.*, viral promoters or leader sequences) can be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding PDE4D, vectors based on SV40 or EBV can be used with an appropriate selectable marker.



### *Bacterial and Yeast Expression Systems*

In bacterial systems, a number of expression vectors can be selected. For example, when a large quantity of PDE4D is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene). In a BLUESCRIPT vector, a sequence encoding PDE4D can be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of  $\beta$ -galactosidase so that a hybrid protein is produced. pIN vectors or pGEX vectors (Promega, Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

### *Plant and Insect Expression Systems*

If plant expression vectors are used, the expression of sequences encoding PDE4D can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used. These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection.

An insect system also can be used to express PDE4D. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia*

- 29 -

larvae. Sequences encoding PDE4D can be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of PDE4D will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which PDE4D can be expressed.

#### *Mammalian Expression Systems*

A number of viral-based expression systems can be used to express PDE4D in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding PDE4D can be ligated into an adenovirus transcription/-translation complex comprising the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing PDE4D in infected host cells [Engelhard, 1994]. If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (e.g., liposomes, polycationic amino polymers, or vesicles). Specific initiation signals also can be used to achieve more efficient translation of sequences encoding PDE4D. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding PDE4D, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert.

- 30 -

Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic.

#### *Host Cells*

5

A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed PDE4D in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the polypeptide also can be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, VA 20110-2209) and can be chosen to ensure the correct modification and processing of the foreign protein.

15

Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express PDE4D can be transformed using expression vectors which can contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 1-2 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced PDE4D sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. Any number of selection systems can be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase [Logan, (1984)] and adenine phosphoribosyltransferase [Wigler, (1977)] genes which can be employed in *tk* or *aprt* cells, respectively. Also, antimetabolite, antibiotic, or herbicide resistance

20

25

30

- 31 -

can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate [Lowy, (1980)], *npt* confers resistance to the aminoglycosides, neomycin and G-418 [Wigler, (1980)], and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively [Colbere-Garapin, 1981].

5 Additional selectable genes have been described. For example, *trpB* allows cells to utilize indole in place of tryptophan, or *hisD*, which allows cells to utilize histinol in place of histidine. Visible markers such as anthocyanins,  $\beta$ -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression

10 attributable to a specific vector system

#### *Detecting Polypeptide Expression*

Although the presence of marker gene expression suggests that a PDE4D polynucleotide is also present, its presence and expression may need to be confirmed.

15 For example, if a sequence encoding PDE4D is inserted within a marker gene sequence, transformed cells containing sequences which encode PDE4D can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding PDE4D under the control of a single

20 promoter. Expression of the marker gene in response to induction or selection usually indicates expression of PDE4D polynucleotide.

Alternatively, host cells which contain a PDE4D polynucleotide and which express PDE4D can be identified by a variety of procedures known to those of skill in the art.

25 These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip-based technologies for the detection and/or quantification of nucleic acid or protein. For example, the presence of a polynucleotide sequence encoding PDE4D can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides

30 encoding PDE4D. Nucleic acid amplification-based assays involve the use of

- 32 -

oligonucleotides selected from sequences encoding PDE4D to detect transformants which contain a PDE4D polynucleotide.

5 A variety of protocols for detecting and measuring the expression of PDE4D, using either polyclonal or monoclonal antibodies specific for the polypeptide, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radio-immunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal antibodies reactive to two non-interfering epitopes on PDE4D can be used, or a competitive binding assay can be  
10 employed.

A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to  
15 polynucleotides encoding PDE4D include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, sequences encoding PDE4D can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes *in vitro* by addition of labeled nucleotides and an appropriate  
20 RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable reporter molecules or labels which can be used for ease of detection include radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles,  
25 and the like.

#### *Expression and Purification of Polypeptides*

Host cells transformed with PDE4D polynucleotides can be cultured under  
30 conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a transformed cell can be secreted or contained

intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing PDE4D polynucleotides can be designed to contain signal sequences which direct secretion of soluble PDE4D through a prokaryotic or eukaryotic cell membrane or which direct  
5 the membrane insertion of membrane-bound PDE4D.

As discussed above, other constructions can be used to join a sequence encoding PDE4D to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but  
10 are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals; protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase  
15 (Invitrogen, San Diego, CA) between the purification domain and PDE4D also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing PDE4D and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography) Maddox,  
20 (1983)], while the enterokinase cleavage site provides a means for purifying PDE4D from the fusion protein [Porath, (1992)].

### *Chemical Synthesis*

25 Sequences encoding PDE4D can be synthesized, in whole or in part, using chemical methods well known in the art. Alternatively, PDE4D itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques. Protein synthesis can either be performed using manual techniques or by automation. Automated synthesis can be achieved,  
30 for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer).

- 34 -

Optionally, fragments of PDE4D can be separately synthesized and combined using chemical methods to produce a full-length molecule.

5 The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography. The composition of a synthetic PDE4D can be confirmed by amino acid analysis or sequencing. Additionally, any portion of the amino acid sequence of PDE4D can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

#### 10 *Production of Altered Polypeptides*

As will be understood by those of skill in the art, it may be advantageous to produce PDE4D polynucleotides possessing non-naturally occurring codons. For example, 15 codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

20 The nucleotide sequences referred to herein can be engineered using methods generally known in the art to alter PDE4D polynucleotides for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides 25 can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

*Antibodies*

Any type of antibody known in the art can be generated to bind specifically to an epitope of PDE4D.

5

“Antibody” as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv, which are capable of binding an epitope of PDE4D. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, e.g., at least 15, 25, or 50 amino acid. An antibody which specifically binds to an epitope of PDE4D can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radioimmunoassays, immunohistochemical assays, immunoprecipitations, or other immunochemical assays known in the art. Various immunoassays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are well known in the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody which specifically binds to the PDE4D immunogen.

10

Typically, an antibody which specifically binds to PDE4D provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immunochemical assay. Preferably, antibodies which specifically bind to PDE4D do not detect other proteins in immunochemical assays and can immunoprecipitate PDE4D from solution.

20

PDE4D can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, PDE4D can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (e.g., aluminum hydroxide), and surface

25

30



- 36 -

active substances (e.g., lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (*bacilli Calmette-Guerin*) and *Corynebacterium parvum* are especially useful.

5

Monoclonal antibodies which specifically bind to PDE4D can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma  
10 technique [Roberge, (1995)].

In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. Monoclonal and  
15 other antibodies also can be "humanized" to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by  
20 replacing residues which differ from those in the human sequences by site directed mutagenesis of individual residues or by grating of entire complementarity determining regions. Antibodies which specifically bind to PDE4D can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. 5,565,332.

25

Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies which specifically bind to PDE4D. Antibodies with related specificity, but of distinct  
30 idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries. Single-chain antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a

- 37 -

template. Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught. A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology.

Antibodies which specifically bind to PDE4D also can be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents. Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, also can be prepared.

Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which PDE4D is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

#### *Antisense Oligonucleotides*

Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 11 nucleotides in length, but can be at least 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense

- 38 -

oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of PDE4D gene products in the cell.

Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamides, carboxymethyl esters, carbonates, and phosphate triesters.

Modifications of PDE4D gene expression can be obtained by designing antisense oligonucleotides which will form duplexes to the control, 5', or regulatory regions of the PDE4D gene. Oligonucleotides derived from the transcription initiation site, *e.g.*, between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature [Nicholls, (1993)]. An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a PDE4D polynucleotide. Antisense oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a PDE4D polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent PDE4D nucleotides, can provide sufficient targeting specificity for PDE4D mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in

- 39 -

length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular PDE4D polynucleotide sequence. Antisense oligonucleotides can be modified without affecting their ability to hybridize to a PDE4D polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art.

### *Ribozymes*

Ribozymes are RNA molecules with catalytic activity [Uhlmann, (1987)]. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide sequences. The coding sequence of a PDE4D polynucleotide can be used to generate ribozymes which will specifically bind to mRNA transcribed from a PDE4D polynucleotide. Methods of designing and constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art. For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target RNA.

Specific ribozyme cleavage sites within a PDE4D RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target RNA containing the cleavage site can be evaluated for secondary structural features which may render the target inoperable. Suitability of candidate PDE4D RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. The nucleotide sequences shown in SEQ ID NO: 1 and its complement provide sources of suitable hybridization region sequences. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease PDE4D expression. Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells (U.S. 5,641,673). Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

## Screening / Screening Assays

### *Regulators*

5 Regulators as used herein, refer to compounds that affect the activity of PDE4D in vivo and/or in vitro. Regulators can be agonists and antagonists of PDE4D polypeptide and can be compounds that exert their effect on the PDE4D activity via the enzymatic activity, expression, post-translational modifications or by other means. Agonists of PDE4D are molecules which, when bound to PDE4D, increase  
10 or prolong the activity of PDE4D. Agonists of PDE4D include proteins, nucleic acids, carbohydrates, small molecules, or any other molecule which activate PDE4D. Antagonists of PDE4D are molecules which, when bound to PDE4D, decrease the amount or the duration of the activity of PDE4D. Antagonists include proteins, nucleic acids, carbohydrates, antibodies, small molecules, or any other molecule  
15 which decrease the activity of PDE4D.

The term "modulate", as it appears herein, refers to a change in the activity of PDE4D polypeptide. For example, modulation may cause an increase or a decrease in enzymatic activity, binding characteristics, or any other biological, functional, or  
20 immunological properties of PDE4D.

As used herein, the terms "specific binding" or "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein recognized by the binding molecule (i.e., the antigenic determinant or epitope). For  
25 example, if an antibody is specific for epitope "A" the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

30

The invention provides methods (also referred to herein as "screening assays") for identifying compounds which can be used for the treatment of hematological and cardiovascular diseases, disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases. The methods  
5 entail the identification of candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other molecules) which bind to PDE4D and/or have a stimulatory or inhibitory effect on the biological activity of PDE4D or its expression and then determining which of these compounds have an effect on symptoms or diseases regarding the hematological and cardiovascular diseases,  
10 disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases in an *in vivo* assay.

Candidate or test compounds or agents which bind to PDE4D and/or have a stimulatory or inhibitory effect on the activity or the expression of PDE4D are  
15 identified either in assays that employ cells which express PDE4D (cell-based assays) or in assays with isolated PDE4D (cell-free assays). The various assays can employ a variety of variants of PDE4D (e.g., full-length PDE4D, a biologically active fragment of PDE4D, or a fusion protein which includes all or a portion of PDE4D). Moreover, PDE4D can be derived from any suitable mammalian species  
20 (e.g., human PDE4D, rat PDE4D or murine PDE4D). The assay can be a binding assay entailing direct or indirect measurement of the binding of a test compound or a known PDE4D ligand to PDE4D. The assay can also be an activity assay entailing direct or indirect measurement of the activity of PDE4D. The assay can also be an expression assay entailing direct or indirect measurement of the expression of  
25 PDE4D mRNA or PDE4D protein. The various screening assays are combined with an *in vivo* assay entailing measuring the effect of the test compound on the symptoms of hematological and cardiovascular diseases, disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases.

The present invention includes biochemical, cell free assays that allow the identification of inhibitors and agonists of PDEs suitable as lead structures for pharmacological drug development. Such assays involve contacting a form of PDE4D (e.g., full-length PDE4D, a biologically active fragment of PDE4D, or a fusion protein comprising all or a portion of PDE4D) with a test compound and determining the ability of the test compound to act as an antagonist (preferably) or an agonist of the enzymatic activity of PDE4D. In one embodiment, the assay includes monitoring the PDE activity of PDE4D by measuring the conversion of either cAMP or cGMP to its nucleoside monophosphate after contacting PDE4D with a test compound.

For example, cAMP and cGMP levels can be measured by the use of the tritium containing compounds  $^3\text{HcAMP}$  and  $^3\text{HcGMP}$  as described in [Hansen, R.S., and Beavo, J.A., PNAS USA1982;79: 2788-92]. To screen a compound pool comprised of a large number of compounds, the microtiter plate-based scintillation proximity assay (SPA) as described in [Bardelle, C. et al. (1999) Anal. Biochem. 275: 148-155] can be applied.

Alternatively, the phosphodiesterase activity of the recombinant protein can be assayed using a commercially available SPA kit (Amersham Pharmacia). The PDE enzyme hydrolyzes cyclic nucleotides, e.g. cAMP and cGMP to their linear counterparts. The SPA assay utilizes the tritiated cyclic nucleotides [ $^3\text{H}$ ]cAMP or [ $^3\text{H}$ ]cGMP, and is based upon the selective interaction of the tritiated non cyclic product with the SPA beads whereas the cyclic substrates are not effectively binding. Radiolabelled product bound to the scintillation beads generates light that can be analyzed in a scintillation counter.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of PDE4D. Such assays can employ full-length PDE4D, a biologically active fragment of PDE4D, or a fusion protein which includes all or a portion of PDE4D. As described in greater detail



- 44 -

below, the test compound can be obtained by any suitable means, e.g., from conventional compound libraries.

Determining the ability of the test compound to modulate the activity of PDE4D can be accomplished, for example, by determining the ability of PDE4D to bind to or interact with a target molecule. The target molecule can be a molecule with which PDE4D binds or interacts with in nature. The target molecule can be a component of a signal transduction pathway which facilitates transduction of an extracellular signal. The target PDE4D molecule can be, for example, a second intracellular protein which has catalytic activity or a protein which facilitates the association of downstream signaling molecules with PDE4D.

Determining the ability of PDE4D to bind to or interact with a target molecule can be accomplished by one of the methods described above for determining direct binding. In one embodiment, determining the ability of a polypeptide of the invention to bind to or interact with a target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (e.g., intracellular  $\text{Ca}^{2+}$ , diacylglycerol,  $\text{IP}_3$ , etc.), detecting catalytic/enzymatic activity of the target on an appropriate substrate, detecting the induction of a reporter gene (e.g., a regulatory element that is responsive to a polypeptide of the invention operably linked to a nucleic acid encoding a detectable marker, e.g., luciferase), or detecting a cellular response.

In various embodiments of the above assay methods of the present invention, it may be desirable to immobilize PDE4D (or a PDE4D target molecule) to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to PDE4D, or interaction of PDE4D with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and

- 45 -

micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase (GST) fusion proteins or glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical; St. Louis, Mo.) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or PDE4D, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components and complex formation is measured either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of binding or activity of PDE4D can be determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either PDE4D or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated polypeptide of the invention or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals; Rockford, Ill.), and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies reactive with PDE4D or target molecules but which do not interfere with binding of the polypeptide of the invention to its target molecule can be derivatized to the wells of the plate, and unbound target or polypeptide of the invention trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with PDE4D or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with PDE4D or target molecule.

- 46 -

Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with PDE4D, or fragments thereof, and washed. Bound PDE4D is then detected by methods well known in the art. Purified PDE4D can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PDE4D specifically compete with a test compound for binding PDE4D. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PDE4D.

The screening assay can also involve monitoring the expression of PDE4D. For example, regulators of expression of PDE4D can be identified in a method in which a cell is contacted with a candidate compound and the expression of PDE4D protein or mRNA in the cell is determined. The level of expression of PDE4D protein or mRNA the presence of the candidate compound is compared to the level of expression of PDE4D protein or mRNA in the absence of the candidate compound. The candidate compound can then be identified as a regulator of expression of PDE4D based on this comparison. For example, when expression of PDE4D protein or mRNA protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of PDE4D protein or mRNA expression. Alternatively, when expression of PDE4D protein or mRNA is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as

an inhibitor of PDE4D protein or mRNA expression. The level of PDE4D protein or mRNA expression in the cells can be determined by methods described below.

### *Binding Assays*

5

For binding assays, the test compound is preferably a small molecule which binds to and occupies the active site of PDE4D polypeptide, thereby making the ligand binding site inaccessible to substrate such that normal biological activity is prevented. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules. Potential ligands which bind to a polypeptide of the invention include, but are not limited to, the natural ligands of known PDE4D PDEs and analogues or derivatives thereof.

In binding assays, either the test compound or the PDE4D polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to PDE4D polypeptide can then be accomplished, for example, by direct counting of radioemmission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product. Alternatively, binding of a test compound to a PDE4D polypeptide can be determined without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a PDE4D polypeptide. A microphysiometer (*e.g.*, Cytosensor™) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound and PDE4D [Haseloff, (1988)].

Determining the ability of a test compound to bind to PDE4D also can be accomplished using a technology such as real-time Bimolecular Interaction Analysis (BIA) [McConnell, (1992); Sjolander, (1991)]. BIA is a technology for studying bio-

- 48 -

specific interactions in real time, without labeling any of the interactants (e.g., BIAcore™). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

5 In yet another aspect of the invention, a PDE4D-like polypeptide can be used as a "bait protein" in a two-hybrid assay or three-hybrid assay [Szabo, (1995); U.S. 5,283,317], to identify other proteins which bind to or interact with PDE4D and modulate its activity.

10 The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding PDE4D can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct a DNA  
15 sequence that encodes an unidentified protein ("prey" or "sample") can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact *in vivo* to form an protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a  
20 reporter gene (e.g., LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with PDE4D.

25 It may be desirable to immobilize either the PDE4D (or polynucleotide) or the test compound to facilitate separation of the bound form from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either the PDE4D-like polypeptide (or polynucleotide) or the test compound can be  
30 bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or

- 49 -

- particles such as beads (including, but not limited to, latex, polystyrene, or glass beads). Any method known in the art can be used to attach PDE4D-like polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to PDE4D (or a polynucleotide encoding for PDE4D) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

In one embodiment, PDE4D is a fusion protein comprising a domain that allows binding of PDE4D to a solid support. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the non-adsorbed PDE4D; the mixture is then incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either PDE4D (or a polynucleotide encoding PDE4D) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated PDE4D (or a polynucleotide encoding biotinylated PDE4D) or test compounds can be prepared from biotin-NHS (N-hydroxysuccinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies which specifically bind to

- 50 -

PDE4D, polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the active site of PDE4D, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

5

Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which specifically bind to PDE4D polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of PDE4D polypeptide, and SDS gel electrophoresis under non-reducing conditions.

10

Screening for test compounds which bind to a PDE4D polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a PDE4D polypeptide or polynucleotide can be used in a cell-based assay system. A PDE4D polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test compound to PDE4D or a polynucleotide encoding PDE4D is determined as described above.

15

#### *Functional Assays*

20

Test compounds can be tested for the ability to increase or decrease PDE4D activity of a PDE4D polypeptide. The PDE4D activity can be measured, for example, using methods described in the specific examples, below. PDE4D activity can be measured after contacting either a purified PDE4D or an intact cell with a test compound. A test compound which decreases PDE4D activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for decreasing PDE4D activity. A test compound which increases PDE4D activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for increasing PDE4D activity.

25

30

*Gene Expression*

In another embodiment, test compounds which increase or decrease PDE4D gene expression are identified. As used herein, the term "correlates with expression of a polynucleotide" indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding PDE4D, by northern analysis or realtime PCR is indicative of the presence of nucleic acids encoding PDE4D in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding PDE4D. The term "microarray", as used herein, refers to an array of distinct polynucleotides or oligonucleotides arrayed on a substrate, such as paper, nylon or any other type of membrane, filter, chip, glass slide, or any other suitable solid support. A PDE4D polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of PDE4D polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of expression of mRNA or polypeptide in the absence of the test compound. The test compound can then be identified as a regulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of the mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

The level of PDE4D mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of PDE4D polynucleotide can be determined, for example, using a variety of techniques known in the art, including immunochemical methods such as radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined *in vivo*, in a cell culture, or in an *in vitro* translation system by detecting incorporation of labelled amino acids into PDE4D.



Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses PDE4D polynucleotide can be used in a cell-based assay system. The PDE4D polynucleotide can be naturally occurring in the cell or  
5 can be introduced using techniques such as those described above. Either a primary culture or an established cell line can be used.

### *Test Compounds*

10 Suitable test compounds for use in the screening assays of the invention can be obtained from any suitable source, e.g., conventional compound libraries. The test compounds can also be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries;  
15 spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of  
20 compounds [Lam, (1997)]. Examples of methods for the synthesis of molecular libraries can be found in the art. Libraries of compounds may be presented in solution or on beads, bacteria, spores, plasmids or phage.

### *Modeling of Regulators*

25 Computer modeling and searching technologies permit identification of compounds, or the improvement of already identified compounds, that can modulate PDE4D expression or activity. Having identified such a compound or composition, the active sites or regions are identified. Such sites might typically be the enzymatic active site, regulator binding sites, or ligand binding sites. The active site can be identified using  
30 methods known in the art including, for example, from the amino acid sequences of peptides, from the nucleotide sequences of nucleic acids, or from study of complexes

of the relevant compound or composition with its natural ligand. In the latter case, chemical or X-ray crystallographic methods can be used to find the active site by finding where on the factor the complexed ligand is found.

5       Next, the three dimensional geometric structure of the active site is determined. This can be done by known methods, including X-ray crystallography, which can determine a complete molecular structure. On the other hand, solid or liquid phase NMR can be used to determine certain intramolecular distances. Any other  
10       experimental method of structure determination can be used to obtain partial or complete geometric structures. The geometric structures may be measured with a complexed ligand, natural or artificial, which may increase the accuracy of the active site structure determined.

15       If an incomplete or insufficiently accurate structure is determined, the methods of computer based numerical modeling can be used to complete the structure or improve its accuracy. Any recognized modeling method may be used, including parameterized models specific to particular biopolymers such as proteins or nucleic acids, molecular dynamics models based on computing molecular motions, statistical  
20       mechanics models based on thermal ensembles, or combined models. For most types of models, standard molecular force fields, representing the forces between constituent atoms and groups, are necessary, and can be selected from force fields known in physical chemistry. The incomplete or less accurate experimental structures can serve as constraints on the complete and more accurate structures  
25       computed by these modeling methods.

30       Finally, having determined the structure of the active site, either experimentally, by modeling, or by a combination, candidate modulating compounds can be identified by searching databases containing compounds along with information on their molecular structure. Such a search seeks compounds having structures that match the determined active site structure and that interact with the groups defining the active

- 54 -

site. Such a search can be manual, but is preferably computer assisted. These compounds found from this search are potential PDE4D modulating compounds.

Alternatively, these methods can be used to identify improved modulating compounds from an already known modulating compound or ligand. The composition of the known compound can be modified and the structural effects of modification can be determined using the experimental and computer modeling methods described above applied to the new composition. The altered structure is then compared to the active site structure of the compound to determine if an improved fit or interaction results. In this manner systematic variations in composition, such as by varying side groups, can be quickly evaluated to obtain modified modulating compounds or ligands of improved specificity or activity.

#### Therapeutic Indications and Methods

It was found by the present applicant that PDE4D is expressed in various human tissues.

#### *Neurology*

CNS disorders include disorders of the central nervous system as well as disorders of the peripheral nervous system.

CNS disorders include, but are not limited to brain injuries, cerebrovascular diseases and their consequences, Parkinson's disease, corticobasal degeneration, motor neuron disease, dementia, including ALS, multiple sclerosis, traumatic brain injury, stroke, post-stroke, post-traumatic brain injury, and small-vessel cerebrovascular disease. Dementias, such as Alzheimer's disease, vascular dementia, dementia with Lewy bodies, frontotemporal dementia and Parkinsonism linked to chromosome 17, frontotemporal dementias, including Pick's disease, progressive nuclear palsy, corticobasal degeneration, Huntington's disease, thalamic degeneration, Creutzfeld-

Jakob dementia, HIV dementia, schizophrenia with dementia, and Korsakoff's psychosis, within the meaning of the definition are also considered to be CNS disorders.

5 Similarly, cognitive-related disorders, such as mild cognitive impairment, age-associated memory impairment, age-related cognitive decline, vascular cognitive impairment, attention deficit disorders, attention deficit hyperactivity disorders, and memory disturbances in children with learning disabilities are also considered to be CNS disorders.

10

Pain, within the meaning of this definition, is also considered to be a CNS disorder. Pain can be associated with CNS disorders, such as multiple sclerosis, spinal cord injury, sciatica, failed back surgery syndrome, traumatic brain injury, epilepsy, Parkinson's disease, post-stroke, and vascular lesions in the brain and spinal cord (e.g., infarct, hemorrhage, vascular malformation). Non-central neuropathic pain includes that associated with post mastectomy pain, phantom feeling, reflex sympathetic dystrophy (RSD), trigeminal neuralgia, radioculopathy, post-surgical pain, HIV/AIDS related pain, cancer pain, metabolic neuropathies (e.g., diabetic neuropathy, vasculitic neuropathy secondary to connective tissue disease),  
15 paraneoplastic polyneuropathy associated, for example, with carcinoma of lung, or leukemia, or lymphoma, or carcinoma of prostate, colon or stomach, trigeminal neuralgia, cranial neuralgias, and post-herpetic neuralgia. Pain associated with peripheral nerve damage, central pain (i.e. due to cerebral ischemia) and various chronic pain i.e., lumbago, back pain (low back pain), inflammatory and/or  
20 rheumatic pain. Headache pain (for example, migraine with aura, migraine without aura, and other migraine disorders), episodic and chronic tension-type headache, tension-type like headache, cluster headache, and chronic paroxysmal hemicrania are also CNS disorders.

25

30 Visceral pain such as pancreatitis, intestinal cystitis, dysmenorrhea, irritable Bowel syndrome, Crohn's disease, biliary colic, ureteral colic, myocardial infarction and

pain syndromes of the pelvic cavity, e.g., vulvodynia, orchialgia, urethral syndrome and protatodynia are also CNS disorders.

5 Also considered to be a disorder of the nervous system are acute pain, for example postoperative pain, and pain after trauma.

The human PDE4D is highly expressed in the following brain tissues: brain, Alzheimer brain, cerebral cortex, Alzheimer cerebral cortex, occipital lobe, parietal lobe, temporal lobe, precentral gyrus, substantia nigra, hippocampus, thalamus, spinal cord. The expression in brain tissues and in particular the differential  
10 expression between diseased tissue Alzheimer brain and healthy tissue brain demonstrates that the human PDE4D or mRNA can be utilized to diagnose nervous system diseases. Additionally the activity of the human PDE4D can be modulated to treat nervous system diseases.

15

#### *Cardiovascular Disorders*

Heart failure is defined as a pathophysiological state in which an abnormality of cardiac function is responsible for the failure of the heart to pump blood at a rate  
20 commensurate with the requirement of the metabolizing tissue. It includes all forms of pumping failures such as high-output and low-output, acute and chronic, right-sided or left-sided, systolic or diastolic, independent of the underlying cause.

Myocardial infarction (MI) is generally caused by an abrupt decrease in coronary  
25 blood flow that follows a thrombotic occlusion of a coronary artery previously narrowed by arteriosclerosis. MI prophylaxis (primary and secondary prevention) is included as well as the acute treatment of MI and the prevention of complications.

Ischemic diseases are conditions in which the coronary flow is restricted resulting in  
30 a perfusion which is inadequate to meet the myocardial requirement for oxygen. This group of diseases includes stable angina, unstable angina and asymptomatic

ischemia.

Arrhythmias include all forms of atrial and ventricular tachyarrhythmias, atrial tachycardia, atrial flutter, atrial fibrillation, atrio-ventricular reentrant tachycardia, preexcitation syndrome, ventricular tachycardia, ventricular flutter, ventricular fibrillation, as well as bradycardic forms of arrhythmias.

Hypertensive vascular diseases include primary as well as all kinds of secondary arterial hypertension, renal, endocrine, neurogenic, others. The genes may be used as drug targets for the treatment of hypertension as well as for the prevention of all complications arising from cardiovascular diseases.

Peripheral vascular diseases are defined as vascular diseases in which arterial and/or venous flow is reduced resulting in an imbalance between blood supply and tissue oxygen demand. It includes chronic peripheral arterial occlusive disease (PAOD), acute arterial thrombosis and embolism, inflammatory vascular disorders, Raynaud's phenomenon and venous disorders.

Atherosclerosis is a cardiovascular disease in which the vessel wall is remodeled, compromising the lumen of the vessel. The atherosclerotic remodeling process involves accumulation of cells, both smooth muscle cells and monocyte/macrophage inflammatory cells, in the intima of the vessel wall. These cells take up lipid, likely from the circulation, to form a mature atherosclerotic lesion. Although the formation of these lesions is a chronic process, occurring over decades of an adult human life, the majority of the morbidity associated with atherosclerosis occurs when a lesion ruptures, releasing thrombogenic debris that rapidly occludes the artery. When such an acute event occurs in the coronary artery, myocardial infarction can ensue, and in the worst case, can result in death.

The formation of the atherosclerotic lesion can be considered to occur in five overlapping stages such as migration, lipid accumulation, recruitment of inflam-

- 58 -

matory cells, proliferation of vascular smooth muscle cells, and extracellular matrix deposition. Each of these processes can be shown to occur in man and in animal models of atherosclerosis, but the relative contribution of each to the pathology and clinical significance of the lesion is unclear.

5

Thus, a need exists for therapeutic methods and agents to treat cardiovascular pathologies, such as atherosclerosis and other conditions related to coronary artery disease.

10

Cardiovascular diseases include but are not limited to disorders of the heart and the vascular system like congestive heart failure, myocardial infarction, ischemic diseases of the heart, all kinds of atrial and ventricular arrhythmias, hypertensive vascular diseases, peripheral vascular diseases, and atherosclerosis.

15

To high or to low levels of fats in the bloodstream, especially cholesterol, can cause long-term problems. The risk to develop atherosclerosis and coronary artery or carotid artery disease (and thus the risk of having a heart attack or stroke) increases with the total cholesterol level increasing. Nevertheless, extremely low cholesterol levels may not be healthy. Examples of disorders of lipid metabolism are

20

hyperlipidemia (abnormally high levels of fats (cholesterol, triglycerides, or both) in the blood, may be caused by family history of hyperlipidemia), obesity, a high-fat diet, lack of exercise, moderate to high alcohol consumption, cigarette smoking, poorly controlled diabetes, and an underactive thyroid gland), hereditary hyperlipidemias (type I hyperlipoproteinemia (familial hyperchylomicronemia), type II hyperlipoproteinemia (familial hypercholesterolemia), type III hyperlipoproteinemia,

25

type IV hyperlipoproteinemia, or type V hyperlipoproteinemia), hypolipoproteinemia, lipidoses (caused by abnormalities in the enzymes that metabolize fats), Gaucher's disease, Niemann-Pick disease, Fabry's disease, Wolman's disease, cerebrotendinous xanthomatosis, sitosterolemia, Refsum's disease, or Tay-Sachs

30

disease.

- 59 -

Kidney disorders may lead to hypertension or hypotension. Examples for kidney problems possibly leading to hypertension are renal artery stenosis, pyelonephritis, glomerulonephritis, kidney tumors, polycystic kidney disease, injury to the kidney, or radiation therapy affecting the kidney. Excessive urination may lead to hypotension.

5

The human PDE4D is highly expressed in the following cardiovascular related tissues: fetal heart, heart, pericardium, heart atrium (right), heart atrium (left), heart ventricle (left), interventricular septum, coronary artery, coronary artery sclerotic. Expression in the above mentioned tissues and in particular the differential expression between diseased tissue coronary artery sclerotic and healthy tissue coronary artery demonstrates that the human PDE4D or mRNA can be utilized to diagnose of cardiovascular diseases. Additionally the activity of the human PDE4D can be modulated to treat cardiovascular diseases.

10

### 15 *Hematological Disorders*

Hematological disorders comprise diseases of the blood and all its constituents as well as diseases of organs and tissues involved in the generation or degradation of all the constituents of the blood. They include but are not limited to 1) Anemias, 2) Myeloproliferative Disorders, 3) Hemorrhagic Disorders, 4) Leukopenia, 5) Eosino-  
philic Disorders, 6) Leukemias, 7) Lymphomas, 8) Plasma Cell Dyscrasias, 9) Disorders of the Spleen in the course of hematological disorders. Disorders according to 1) include, but are not limited to anemias due to defective or deficient hem  
synthesis, deficient erythropoiesis. Disorders according to 2) include, but are not  
limited to polycythemia vera, tumor-associated erythrocytosis, myelofibrosis,  
thrombocythemia. Disorders according to 3) include, but are not limited to vasculitis,  
thrombocytopenia, heparin-induced thrombocytopenia, thrombotic thrombocytopenic  
purpura, hemolytic-uremic syndrome, hereditary and acquired disorders of platelet  
function, hereditary coagulation disorders. Disorders according to 4) include, but are  
not limited to neutropenia, lymphocytopenia. Disorders according to 5) include, but  
are not limited to hypereosinophilia, idiopathic hypereosinophilic syndrome.

20

25

30



- 60 -

Disorders according to 6) include, but are not limited to acute myeloic leukemia, acute lymphoblastic leukemia, chronic myelocytic leukemia, chronic lymphocytic leukemia, myelodysplastic syndrome. Disorders according to 7) include, but are not limited to Hodgkin's disease, non-Hodgkin's lymphoma, Burkitt's lymphoma, mycosis fungoides cutaneous T-cell lymphoma. Disorders according to 8) include, but are not limited to multiple myeloma, macroglobulinemia, heavy chain diseases. In extension of the preceding idiopathic thrombocytopenic purpura, iron deficiency anemia, megaloblastic anemia (vitamin B12 deficiency), aplastic anemia, thalassemia, malignant lymphoma bone marrow invasion, malignant lymphoma skin invasion, hemolytic uremic syndrome, giant platelet disease are considered to be hematological diseases too.

The human PDE4D is highly expressed in the following tissues of the hematological system: leukocytes (peripheral blood), bone marrow, thymus, thrombocytes, bone marrow stromal cells, spleen, spleen liver cirrhosis. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue spleen liver cirrhosis and healthy tissue spleen demonstrates that the human PDE4D or mRNA can be utilized to diagnose of hematological diseases. Additionally the activity of the human PDE4D can be modulated to treat hematological disorders.

#### *Disorders Related to Pulmology*

Asthma is thought to arise as a result of interactions between multiple genetic and environmental factors and is characterized by three major features: 1) intermittent and reversible airway obstruction caused by bronchoconstriction, increased mucus production, and thickening of the walls of the airways that leads to a narrowing of the airways, 2) airway hyperresponsiveness, and 3) airway inflammation. Certain cells are critical to the inflammatory reaction of asthma and they include T cells and antigen presenting cells, B cells that produce IgE, and mast cells, basophils, eosinophils, and other cells that bind IgE. These effector cells accumulate at the site

- 61 -

of allergic reaction in the airways and release toxic products that contribute to the acute pathology and eventually to tissue destruction related to the disorder. Other resident cells, such as smooth muscle cells, lung epithelial cells, mucus-producing cells, and nerve cells may also be abnormal in individuals with asthma and may  
5 contribute to its pathology. While the airway obstruction of asthma, presenting clinically as an intermittent wheeze and shortness of breath, is generally the most pressing symptom of the disease requiring immediate treatment, the inflammation and tissue destruction associated with the disease can lead to irreversible changes that eventually make asthma a chronic and disabling disorder requiring long-term  
10 management.

Chronic obstructive pulmonary (or airways) disease (COPD) is a condition defined physiologically as airflow obstruction that generally results from a mixture of emphysema and peripheral airway obstruction due to chronic bronchitis [Botstein,  
15 1980]. Emphysema is characterised by destruction of alveolar walls leading to abnormal enlargement of the air spaces of the lung. Chronic bronchitis is defined clinically as the presence of chronic productive cough for three months in each of two successive years. In COPD, airflow obstruction is usually progressive and is only partially reversible. By far the most important risk factor for development of  
20 COPD is cigarette smoking, although the disease does also occur in non-smokers.

The human PDE4D is highly expressed in the following tissues of the respiratory system: fetal lung, lung, lung right upper lobe, lung right mid lobe, lung right lower lobe, lung tumor, lung COPD, trachea. The expression in the above mentioned  
25 tissues and in particular the differential expression between diseased tissue lung tumor and healthy tissue lung demonstrates that the human PDE4D or mRNA can be utilized to diagnose of respiratory diseases. Additionally the activity of the human PDE4D can be modulated to treat those diseases.

*Disorders Related to Urology*

Genitourinary disorders comprise benign and malign disorders of the organs constituting the genitourinary system of female and male, renal diseases like acute or chronic renal failure, immunologically mediated renal diseases like renal transplant rejection, lupus nephritis, immune complex renal diseases, glomerulopathies, nephritis, toxic nephropathy, obstructive uropathies like benign prostatic hyperplasia (BPH), neurogenic bladder syndrome, urinary incontinence like urge-, stress-, or overflow incontinence, pelvic pain, and erectile dysfunction.

The human PDE4D is highly expressed in the following urological tissues: prostata, bladder, fetal kidney, kidney, kidney tumor, HEK 293 cells. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue kidney tumor and healthy tissue kidney demonstrates that the human PDE4D or mRNA can be utilized to diagnose of urological disorders. Additionally the activity of the human PDE4D can be modulated to treat urological disorders.

*Applications*

The present invention provides for both prophylactic and therapeutic methods for disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders.

The regulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of PDE4D. An agent that modulates activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of the polypeptide, a peptide, a peptidomimetic, or any small molecule. In one embodiment, the agent stimulates one or more of the biological activities of PDE4D. Examples of such stimulatory agents include the active PDE4D and nucleic acid molecules encoding a portion of PDE4D. In another embodiment, the agent inhibits one or more of the biological activities of PDE4D.

- 63 -

Examples of such inhibitory agents include antisense nucleic acid molecules and antibodies. These regulatory methods can be performed in vitro (e.g., by culturing the cell with the agent) or, alternatively, in vivo (e.g, by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by unwanted expression or activity of PDE4D or a protein in the PDE4D signaling pathway. In one embodiment, the method involves administering an agent like any agent identified or being identifiable by a screening assay as described herein, or combination of such agents that modulate say upregulate or downregulate the expression or activity of PDE4D or of any protein in the PDE4D signaling pathway. In another embodiment, the method involves administering a regulator of PDE4D as therapy to compensate for reduced or undesirably low expression or activity of PDE4D or a protein in the PDE4D signaling pathway.

Stimulation of activity or expression of PDE4D is desirable in situations in which enzymatic activity or expression is abnormally low and in which increased activity is likely to have a beneficial effect. Conversely, inhibition of enzymatic activity or expression of PDE4D is desirable in situations in which activity or expression of PDE4D is abnormally high and in which decreasing its activity is likely to have a beneficial effect.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference.

### Pharmaceutical Compositions

This invention further pertains to novel agents identified by the above-described screening assays and uses thereof for treatments as described herein.

- 64 -

The nucleic acid molecules, polypeptides, and antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier.

5 As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any

10 conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

The invention includes pharmaceutical compositions comprising a regulator of

15 PDE4D expression or activity (and/or a regulator of the activity or expression of a protein in the PDE4D signaling pathway) as well as methods for preparing such compositions by combining one or more such regulators and a pharmaceutically acceptable carrier. Also within the invention are pharmaceutical compositions comprising a regulator identified using the screening assays of the invention

20 packaged with instructions for use. For regulators that are antagonists of PDE4D activity or which reduce PDE4D expression, the instructions would specify use of the pharmaceutical composition for treatment of hematological and cardiovascular diseases, disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases. For regulators that are

25 agonists of PDE4D activity or increase PDE4D expression, the instructions would specify use of the pharmaceutical composition for treatment of hematological and cardiovascular diseases, disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases.

30 An inhibitor of PDE4D may be produced using methods which are generally known in the art. In particular, purified PDE4D may be used to produce antibodies or to

- 65 -

screen libraries of pharmaceutical agents to identify those which specifically bind PDE4D. Antibodies to PDE4D may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain antibodies, Fab fragments, and fragments  
5 produced by a Fab expression library. Neutralizing antibodies like those which inhibit dimer formation are especially preferred for therapeutic use.

In another embodiment of the invention, the polynucleotides encoding PDE4D, or any fragment or complement thereof, may be used for therapeutic purposes. In one  
10 aspect, the complement of the polynucleotide encoding PDE4D may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding PDE4D. Thus, complementary molecules or fragments may be used to modulate PDE4D activity, or to achieve regulation of gene function. Such  
15 technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding PDE4D.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia  
20 viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors which will express nucleic acid sequence complementary to the polynucleotides of the gene encoding PDE4D. These techniques are described, for example, in [Scott and Smith (1990)].

25 Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

30 An additional embodiment of the invention relates to the administration of a pharmaceutical composition containing PDE4D in conjunction with a pharma-

ceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of PDE4D, antibodies to PDE4D, and mimetics, agonists, antagonists, or inhibitors of PDE4D. The compositions may be administered alone or in combination with at least one other agent, such as a  
5 stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

10 A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include  
15 the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and  
20 agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

25 Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EM<sup>TM</sup> (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In  
30 all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and

- 67 -

storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, a pharmaceutically acceptable polyol like glycerol, propylene glycol, liquid polyethylene glycol, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin. Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a polypeptide or antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed.



- 68 -

Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient  
5 such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

10 For administration by inhalation, the compounds are delivered in the form of an aerosol spray from a pressurized container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For  
15 transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the  
20 active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with  
25 conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect  
the compound against rapid elimination from the body, such as a controlled release  
formulation, including implants and microencapsulated delivery systems. Bio-  
30 degradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid.

- 69 -

Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as  
5 pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as  
10 used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active  
15 compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration. For pharmaceutical compositions which include an antagonist of PDE4D activity, a compound which reduces expression of PDE4D, or a compound which reduces expression or activity of a protein in the PDE4D signaling pathway or any combination thereof, the instructions for administration will specify use of the composition for hematological and  
20 cardiovascular diseases, disorders of the peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases. For pharmaceutical compositions which include an agonist of PDE4D activity, a compound which increases expression of PDE4D, or a compound which increases expression or activity of a protein in the PDE4D signaling pathway or any  
25 combination thereof, the instructions for administration will specify use of the composition for hematological and cardiovascular diseases, disorders of the  
30

- 70 -

peripheral and central nervous system, COPD, asthma, genito-urological disorders and inflammation diseases.

### Diagnostics

5

In another embodiment, antibodies which specifically bind PDE4D may be used for the diagnosis of disorders characterized by the expression of PDE4D, or in assays to monitor patients being treated with PDE4D or agonists, antagonists, and inhibitors of PDE4D. Antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for PDE4D include methods which utilize the antibody and a label to detect PDE4D in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent joining with a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

15

A variety of protocols for measuring PDE4D, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of PDE4D expression. Normal or standard values for PDE4D expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to PDE4D under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, preferably by photometric means. Quantities of PDE4D expressed in subject samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

20

25

In another embodiment of the invention, the polynucleotides encoding PDE4D may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied

30

- 71 -

tissues in which expression of PDE4D may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of PDE4D, and to monitor regulation of PDE4D levels during therapeutic intervention.

5

Polynucleotide sequences encoding PDE4D may be used for the diagnosis of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders associated with expression of PDE4D. The polynucleotide sequences encoding PDE4D may be used  
10 in Southern, Northern, or dot-blot analysis, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; and in microarrays utilizing fluids or tissues from patient biopsies to detect altered PDE4D expression. Such qualitative or quantitative methods are well known in the art.

15 In a particular aspect, the nucleotide sequences encoding PDE4D may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding PDE4D may be labelled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation  
20 period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly altered from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding PDE4D in the sample indicates the presence of the  
25 associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

30 In order to provide a basis for the diagnosis of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders associated with expression of PDE4D, a normal or

standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding PDE4D, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

#### **Determination of a Therapeutically Effective Dose**

The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases PDE4D activity relative to PDE4D activity which occurs in the absence of the therapeutically effective dose. For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

Therapeutic efficacy and toxicity, e.g.,  $ED_{50}$  (the dose therapeutically effective in 50% of the population) and  $LD_{50}$  (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio,  $LD_{50}/ED_{50}$ . Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the  $ED_{50}$  with little or no toxicity. The dosage varies

- 73 -

within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration. The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors which can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts can vary from 0.1 micrograms to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc. If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either *ex vivo* or *in vivo* using well-established techniques including, but not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, "gene gun", and DEAE- or calcium phosphate-mediated transfection.

If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a ribozyme. Polynucleotides which express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described above. Preferably, a reagent reduces expression of PDE4D gene or the activity of PDE4D by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism

- 74 -

chosen to decrease the level of expression of PDE4D gene or the activity of PDE4D can be assessed using methods well known in the art, such as hybridization of nucleotide probes to PDE4D-specific mRNA, quantitative RT-PCR, immunologic detection of PDE4D, or measurement of PDE4D activity.

5

In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects. Any of the therapeutic methods described above can be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

10  
15

Nucleic acid molecules of the invention are those nucleic acid molecules which are contained in a group of nucleic acid molecules consisting of (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2, (ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1, (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1, (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii); and (v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code, wherein the polypeptide encoded by said nucleic acid molecule has PDE4D activity.

20  
25

Polypeptides of the invention are those polypeptides which are contained in a group of polypeptides consisting of (i) polypeptides having the sequence of SEQ ID NO: 2, (ii) polypeptides comprising the sequence of SEQ ID NO: 2, (iii) polypeptides

30

- encoded by nucleic acid molecules of the invention and (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% homology with a polypeptide of (i), (ii), or (iii), wherein said purified polypeptide has PDE4D activity.

5      An object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising the steps of (i) contacting a test compound with a PDE4D polypeptide, (ii) detect binding  
10      of said test compound to said PDE4D polypeptide. E.g., compounds that bind to the PDE4D polypeptide are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders  
15      of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising the steps of (i) determining the activity of a PDE4D polypeptide at a certain concentration of a test compound or in the absence of said test compound, (ii) determining the activity of said polypeptide at a different concentration of said test compound.  
20      E.g., compounds that lead to a difference in the activity of the PDE4D polypeptide in (i) and (ii) are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders  
25      of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising the steps of (i) determining the activity of a PDE4D polypeptide at a certain concentration of a test compound, (ii) determining the activity of a PDE4D polypeptide at the presence of a compound known to be a regulator of a PDE4D  
30      polypeptide. E.g., compounds that show similar effects on the activity of the PDE4D



- 76 -

polypeptide in (i) as compared to compounds used in (ii) are identified potential therapeutic agents for such a disease.

5 Other objects of the invention are methods of the above, wherein the step of contacting is in or at the surface of a cell.

Other objects of the invention are methods of the above, wherein the cell is in vitro.

10 Other objects of the invention are methods of the above, wherein the step of contacting is in a cell-free system.

Other objects of the invention are methods of the above, wherein the polypeptide is coupled to a detectable label.

15 Other objects of the invention are methods of the above, wherein the compound is coupled to a detectable label.

Other objects of the invention are methods of the above, wherein the test compound displaces a ligand which is first bound to the polypeptide.

20 Other objects of the invention are methods of the above, wherein the polypeptide is attached to a solid support.

25 Other objects of the invention are methods of the above, wherein the compound is attached to a solid support.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological  
30 diseases, respiratory diseases and urological disorders in a mammal comprising the steps of (i) contacting a test compound with a PDE4D polynucleotide, (ii) detect

- 77 -

binding of said test compound to said PDE4D polynucleotide. Compounds that, e.g., bind to the PDE4D polynucleotide are potential therapeutic agents for the treatment of such diseases.

- 5 Another object of the invention is the method of the above, wherein the nucleic acid molecule is RNA.

Another object of the invention is a method of the above, wherein the contacting step is in or at the surface of a cell.

10

Another object of the invention is a method of the above, wherein the contacting step is in a cell-free system.

15

Another object of the invention is a method of the above, wherein the polynucleotide is coupled to a detectable label.

Another object of the invention is a method of the above, wherein the test compound is coupled to a detectable label.

20

Another object of the invention is a method of diagnosing a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising the steps of (i) determining the amount of a PDE4D polynucleotide in a sample taken from said mammal, (ii) determining the amount of PDE4D polynucleotide in healthy and/or diseased mammal. A disease is diagnosed, e.g., if there is a substantial similarity in the amount of PDE4D polynucleotide in said test mammal as compared to a diseased mammal.

25

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases,

30

- 78 -

respiratory diseases and urological disorders in a mammal comprising a therapeutic agent which binds to a PDE4D polypeptide.

5 Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising a therapeutic agent which regulates the activity of a PDE4D polypeptide.

10 Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising a therapeutic agent which regulates the activity of a PDE4D polypeptide, wherein said therapeutic  
15 agent is (i) a small molecule, (ii) an RNA molecule, (iii) an antisense oligonucleotide, (iv) a polypeptide, (v) an antibody, or (vi) a ribozyme.

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral  
20 and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising a PDE4D polynucleotide.

25 Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising a PDE4D polypeptide.

30 Another object of the invention is the use of regulators of a PDE4D for the preparation of a pharmaceutical composition for the treatment of a disease comprised

- 79 -

in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal.

5 Another object of the invention is a method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising the steps of (i) identifying a regulator of PDE4D, (ii)  
10 determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal; and (iii) combining of said regulator with an acceptable pharmaceutical carrier.

15 Another object of the invention is the use of a regulator of PDE4D for the regulation of PDE4D activity in a mammal having a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders.

20 The examples below are provided to illustrate the subject invention. These examples are provided by way of illustration and are not included for the purpose of limiting the invention.

- 80 -

Examples**Example 1: Search for homologous sequences in public sequence data bases**

5 The degree of homology can readily be calculated by known methods. Preferred methods to determine homology are designed to give the largest match between the sequences tested. Methods to determine homology are codified in publicly available computer programs such as BestFit, BLASTP, BLASTN, and FASTA. The BLAST programs are publicly available from NCBI and other sources in the internet.

10

For PDE4D the following hits to known sequences were identified by using the BLAST algorithm [Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ; Nucleic Acids Res 1997 Sep 1; 25(17): 3389-402] and the following set of parameters: matrix = BLOSUM62 and low complexity filter. The following  
15 databases were searched: NCBI (non-redundant database) and DERWENT patent database (Geneseq).

The following hits were found:

20 >ref|NM\_006203.1| Homo sapiens phosphodiesterase 4D, cAMP-specific (phosphodiesterase E3 dunce homolog, Drosophila) (PDE4D), mRNA, Length = 3829, Score = 7352 bits (3823), Expect = 0.0, Identities = 3829/3829 (100%)

25 >gb|L20969.1|HUMPDEE Homo sapiens cyclic AMP phosphodiesterase mRNA, complete cds, Length = 3829, Score = 7352 bits (3823), Expect = 0.0, Identities = 3829/3829 (100%)

30 >NA2001B:ABV24957 Abv24957 Human prostate expression marker cDNA, 24948. 9/2002, Length = 7146, Score = 7331 bits (3812), Expect = 0.0, Identities = 3825/3828 (99%), Gaps = 1/3828 (0%)

35 >NA2001B:AAS71151 Aas71151 DNA encoding novel human diagnostic protein #6955. 2/2002, Length = 3130, Score = 5679 bits (2953), Expect = 0.0, Identities = 2953/2953 (100%)

- 81 -

- 5 >NA2000:AAA88199 Aaa88199 pPDE43 human dunce-like PDE nucleotide  
sequence SEQ ID NO:62. 12/2000, Length = 3130, Score = 5679 bits  
(2953), Expect = 0.0, Identities = 2953/2953 (100%)
- >NA1999:AAZ32264 Aaz32264 Human dunce-like phosphodiesterase pPDE43,  
encoding cDNA. 1/2000, Length = 3130, Score = 5679 bits (2953),  
Expect = 0.0, Identities = 2953/2953 (100%)
- 10 >gb|AR083672.1|AR083672 Sequence 62 from patent US 5977305, Length =  
3130, Score = 5679 bits (2953), Expect = 0.0, Identities = 2953/2953  
(100%)
- 15 >gb|AF012073.1|AF012073 Homo sapiens cAMP-specific phosphodiesterase  
PDE4D5 (PDE4D) mRNA, complete cds, Length = 3332, Score = 5679 bits  
(2953), Expect = 0.0, Identities = 2953/2953 (100%)
- 20 >gb|L20970.1|HUMPDEF Human phosphodiesterase mRNA, complete cds,  
Length = 3130, Score = 5679 bits (2953), Expect = 0.0, Identities =  
2953/2953 (100%)
- 25 >ref|XM\_056815.5| Homo sapiens phosphodiesterase 4D, cAMP-specific  
(phosphodiesterase E3 dunce homolog, Drosophila) (PDE4D), mRNA,  
Length = 5967; Score = 5664 bits (2945), Expect = 0.0, Identities =  
2952/2953 (99%), Gaps = 1/2953 (0%)
- 30 >NA2002:ABK84184 Abk84184 Human cDNA differentially expressed in  
granulocytic cells #755. 8/2002, Length = 5706, Score = 5502 bits  
(2861), Expect = 0.0, Identities = 2874/2878 (99%), Gaps = 1/2878  
(0%)
- 35 >gb|U02882.1|HSU02882 Human rolipram-sensitive 3',5'-cyclic AMP  
phosphodiesterase mRNA, complete cds, Length = 5706, Score = 5502  
bits (2861), Expect = 0.0, Identities = 2874/2878 (99%), Gaps =  
1/2878 (0%)
- >gb|AF012074.1|AF012074 Homo sapiens cAMP-specific phosphodiesterase  
PDE4D2 (PDE4D) mRNA, complete cds, Length = 2984, Score = 4839 bits  
(2516), Expect = 0.0, Identities = 2516/2516 (100%)

**Example 2: Expression profiling**

5 Total cellular RNA was isolated from cells by one of two standard methods: 1) guanidine isothiocyanate/Cesium chloride density gradient centrifugation [Kellogg, (1990)]; or with the Tri-Reagent protocol according to the manufacturer's specifications (Molecular Research Center, Inc., Cincinnati, Ohio). Total RNA prepared by the Tri-reagent protocol was treated with DNase I to remove genomic DNA contamination.

10 For relative quantitation of the mRNA distribution of PDE4D, total RNA from each cell or tissue source was first reverse transcribed. 85 µg of total RNA was reverse transcribed using 1 µmole random hexamer primers, 0.5 mM each of dATP, dCTP, dGTP and dTTP (Qiagen, Hilden, Germany), 3000 U RnaseQut (Invitrogen, Groningen, Netherlands) in a final volume of 680 µl. The first strand synthesis buffer and Omniscript reverse transcriptase (2 u/µl) were from (Qiagen, Hilden, Germany). The reaction was incubated at 37°C for 90 minutes and cooled on ice. The volume was adjusted to 6800 µl with water, yielding a final concentration of 12.5 ng/µl of starting RNA.

20 For relative quantitation of the distribution of PDE4D mRNA in cells and tissues the Perkin Elmer ABI Prism RTM. 7700 Sequence Detection system or Biorad iCycler was used according to the manufacturer's specifications and protocols. PCR reactions were set up to quantitate PDE4D and the housekeeping genes HPRT (hypoxanthine phosphoribosyltransferase), GAPDH (glyceraldehyde-3-phosphate dehydrogenase), β-actin, and others. Forward and reverse primers and probes for PDE4D were designed using the Perkin Elmer ABI Primer Express™ software and were synthesized by TibMolBiol (Berlin, Germany). The PDE4D forward primer sequence was: Primer1 (SEQ ID NO: 3). The PDE4D reverse primer sequence was  
25 Primer2 (SEQ ID NO: 5). Probe1 (SEQ ID NO: 4), labelled with FAM (carboxy-fluorescein succinimidyl ester) as the reporter dye and TAMRA (carboxytetra-  
30

methyrlrhodamine) as the quencher, is used as a probe for PDE4D. The following reagents were prepared in a total of 25  $\mu$ l : 1x TaqMan buffer A, 5.5 mM  $MgCl_2$ , 200 nM of dATP, dCTP, dGTP, and dUTP, 0.025 U/ $\mu$ l AmpliTaq Gold<sup>TM</sup>, 0.01 U/ $\mu$ l AmpErase and Probe1 (SEQ ID NO: 4), PDE4D forward and reverse primers each at 200 nM, 200 nM PDE4D FAM/TAMRA-labelled probe, and 5  $\mu$ l of template cDNA. Thermal cycling parameters were 2 min at 50°C, followed by 10 min at 95°C, followed by 40 cycles of melting at 95°C for 15 sec and annealing/extending at 60°C for 1 min.

#### 10 *Calculation of corrected CT values*

The CT (threshold cycle) value is calculated as described in the "Quantitative determination of nucleic acids" section. The CF-value (factor for threshold cycle correction) is calculated as follows :

15

1. PCR reactions were set up to quantitate the housekeeping genes (HKG) for each cDNA sample.

20

2.  $CT_{HKG}$ -values (threshold cycle for housekeeping gene) were calculated as described in the "Quantitative determination of nucleic acids" section.

3.  $CT_{HKG}$ -mean values (CT mean value of all HKG tested on one cDNAs) of all HKG for each cDNA are calculated ( $n$  = number of HKG):

25

$$CT_{HKG-n}\text{-mean value} = (CT_{HKG1}\text{-value} + CT_{HKG2}\text{-value} + \dots + CT_{HKG-n}\text{-value}) / n$$

4.  $CT_{\text{panel}}$  mean value (CT mean value of all HKG in all tested cDNAs) =

$$(CT_{HKG1}\text{-mean value} + CT_{HKG2}\text{-mean value} + \dots + CT_{HKG-y}\text{-mean value}) / y$$

30

( $y$  = number of cDNAs)



- 84 -

5.  $CF_{cDNA-n}$  (correction factor for cDNA n) =  $CT_{panel-mean\ value} - CT_{HKG-n-mean\ value}$

6.  $CT_{cDNA-n}$  (CT value of the tested gene for the cDNA n) +  $CF_{cDNA-n}$  (correction  
5 factor for cDNA n) =  $CT_{cor-cDNA-n}$  (corrected CT value for a gene on cDNA n)

### *Calculation of relative expression*

Definition : highest  $CT_{cor-cDNA-n} \neq 40$  is defined as  $CT_{cor-cDNA [high]}$

10 Relative Expression =  $2^{(CT_{cor-cDNA [high]} - CT_{cor-cDNA-n})}$

### *Tissues*

The expression of PDE4D was investigated in the tissues listed in table 1.

15

### *Expression profile*

The results of the the mRNA-quantification (expression profiling) is shown in  
Table 1.

20

Table 1: *Relative expression of PDE4D in various human tissues*

<u>Tissue</u>	<u>Relative Expression</u>
fetal heart	129
25 heart	83
pericardium	71
heart atrium (right)	26
heart atrium (left)	138
heart ventricle (left)	20
30 interventricular septum	33
fetal aorta	1
aorta	2

- 85 -

	aorta sclerotic	2
	artery	5
	coronary artery	15
	coronary artery sclerotic	5
5	vein	6
	coronary artery smooth muscle primary cells	6
	HUVEC cells	54
	skin	38
10	adrenal gland	2
	thyroid	56
	thyroid tumor	21
	pancreas	2
15	pancreas liver cirrhosis	18
	esophagus	53
	esophagus tumor	24
	stomach	15
20	stomach tumor	68
	colon	30
	colon tumor	9
	small intestine	8
	ileum	62
25	ileum tumor	2
	ileum chronic inflammation	7
	rectum	298
	salivary gland	10
	fetal liver	4
30	liver	24
	liver liver cirrhosis	76
	liver tumor	38
	HEP G2 cells	39

- 86 -

	leukocytes (peripheral blood)	88
	Jurkat (T-cells)	2
	bone marrow	14
	erythrocytes	4
5	lymphnode	5
	thymus	20
	thrombocytes	16
	bone marrow stromal cells	16
	bone marrow CD71+ cells	2
10	bone marrow CD33+ cells	5
	bone marrow CD34+ cells	5
	bone marrow CD15+ cells	4
	cord blood CD71+ cells	2
	spleen	38
15	spleen liver cirrhosis	8
	skeletal muscle	272
	adipose	22
20	fetal brain	152
	brain	22
	Alzheimer brain	47
	cerebellum	1
	cerebellum (right)	29
25	cerebellum (left)	21
	cerebral cortex	42
	Alzheimer cerebral cortex	204
	frontal lobe	25
	Alzheimer brain frontal lobe	32
30	occipital lobe	181
	parietal lobe	182
	temporal lobe	237
	precentral gyrus	199
	postcentral gyrus	6

- 87 -

	tonsilla cerebelli	18
	vermis cerebelli	10
	pons	13
	substantia nigra	109
5	cerebral meninges	3
	cerebral peduncles	30
	corpus callosum	14
	hippocampus	218
	thalamus	62
10	dorsal root ganglia	16
	spinal cord	42
	neuroblastoma SK-N-MC cells	29
	neuroblastoma SH-SY5Y cells	16
	neuroblastoma IMR32 cells	7
15	glial tumor H4 cells	28
	retina	26
	fetal lung	104
	fetal lung fibroblast IMR-90 cells	8
20	lung	16
	lung right upper lobe	49
	lung right mid lobe	25
	lung right lower lobe	23
	lung tumor	43
25	lung COPD	8
	trachea	10
	cervix	12
	testis	9
30	HeLa cells (cervix tumor)	26
	placenta	25
	uterus	29
	uterus tumor	21
	ovary	21

- 88 -

	ovary tumor	10
	breast	39
	breast tumor	18
	MDA MB 231 cells (breast tumor)	17
5	mammary gland	6
	prostate	20
	prostate BPH	6
	bladder	140
10	ureter	22
	penis	12
	corpus cavernosum	10
	fetal kidney	284
	kidney	48
15	kidney tumor	20
	HEK 293 cells	38

### Example 3: Antisense Analysis

20 Knowledge of the correct, complete cDNA sequence coding for PDE4D enables its use as a tool for antisense technology in the investigation of gene function. Oligonucleotides, cDNA or genomic fragments comprising the antisense strand of a polynucleotide coding for PDE4D are used either in vitro or in vivo to inhibit translation of the mRNA. Such technology is now well known in the art, and

25 antisense molecules can be designed at various locations along the nucleotide sequences. By treatment of cells or whole test animals with such antisense sequences, the gene of interest is effectively turned off. Frequently, the function of the gene is ascertained by observing behavior at the intracellular, cellular, tissue or organismal level (e.g., lethality, loss of differentiated function, changes in

30 morphology, etc.).

In addition to using sequences constructed to interrupt transcription of a particular open reading frame, modifications of gene expression is obtained by designing antisense sequences to intron regions, promoter/enhancer elements, or even to trans-acting regulatory genes.

5

#### Example 4: Expression of PDE4D

Expression of PDE4D is accomplished by subcloning the cDNAs into appropriate expression vectors and transfecting the vectors into expression hosts such as, e.g., *E. coli*. In a particular case, the vector is engineered such that it contains a promoter for  $\beta$ -galactosidase, upstream of the cloning site, followed by sequence containing the amino-terminal Methionine and the subsequent seven residues of  $\beta$ -galactosidase. Immediately following these eight residues is an engineered bacteriophage promoter useful for artificial priming and transcription and for providing a number of unique endonuclease restriction sites for cloning.

15

Induction of the isolated, transfected bacterial strain with Isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) using standard methods produces a fusion protein corresponding to the first seven residues of  $\beta$ -galactosidase, about 15 residues of "linker", and the peptide encoded within the cDNA. Since cDNA clone inserts are generated by an essentially random process, there is probability of 33% that the included cDNA will lie in the correct reading frame for proper translation. If the cDNA is not in the proper reading frame, it is obtained by deletion or insertion of the appropriate number of bases using well known methods including in vitro mutagenesis, digestion with exonuclease III or mung bean nuclease, or the inclusion of an oligonucleotide linker of appropriate length.

20

25

The PDE4D cDNA is shuttled into other vectors known to be useful for expression of proteins in specific hosts. Oligonucleotide primers containing cloning sites as well as a segment of DNA (about 25 bases) sufficient to hybridize to stretches at both ends of the target cDNA is synthesized chemically by standard methods. These

30

- 90 -

primers are then used to amplify the desired gene segment by PCR. The resulting gene segment is digested with appropriate restriction enzymes under standard conditions and isolated by gel electrophoresis. Alternately, similar gene segments are produced by digestion of the cDNA with appropriate restriction enzymes. Using appropriate primers, segments of coding sequence from more than one gene are ligated together and cloned in appropriate vectors. It is possible to optimize expression by construction of such chimeric sequences.

Suitable expression hosts for such chimeric molecules include, but are not limited to, mammalian cells such as Chinese Hamster Ovary (CHO) and human 293 cells., insect cells such as Sf9 cells, yeast cells such as *Saccharomyces cerevisiae* and bacterial cells such as *E. coli*. For each of these cell systems, a useful expression vector also includes an origin of replication to allow propagation in bacteria, and a selectable marker such as the  $\beta$ -lactamase antibiotic resistance gene to allow plasmid selection in bacteria. In addition, the vector may include a second selectable marker such as the neomycin phosphotransferase gene to allow selection in transfected eukaryotic host cells. Vectors for use in eukaryotic expression hosts require RNA processing elements such as 3' polyadenylation sequences if such are not part of the cDNA of interest.

Additionally, the vector contains promoters or enhancers which increase gene expression. Such promoters are host specific and include MMTV, SV40, and metallothionine promoters for CHO cells; trp, lac, tac and T7 promoters for bacterial hosts; and alpha factor, alcohol oxidase and PGH promoters for yeast. Transcription enhancers, such as the rous sarcoma virus enhancer, are used in mammalian host cells. Once homogeneous cultures of recombinant cells are obtained through standard culture methods, large quantities of recombinantly produced PDE4D are recovered from the conditioned medium and analyzed using chromatographic methods known in the art. For example, PDE4D can be cloned into the expression vector pcDNA3, as exemplified herein. This product can be used to transform, for example, HEK293 or COS by methodology standard in the art. Specifically, for

example, using Lipofectamine (Gibco BRL catalog no. 18324-020) mediated gene transfer.

#### Example 5: Isolation of Recombinant PDE4D

5

PDE4D is expressed as a chimeric protein with one or more additional polypeptide domains added to facilitate protein purification. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals [Appa Rao, 1997] and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Washington). The inclusion of a cleavable linker sequence such as Factor Xa or enterokinase (Invitrogen, Groningen, The Netherlands) between the purification domain and the PDE4D sequence is useful to facilitate expression of PDE4D.

15

The following example provides a method for purifying PDE4D.

20

PDE4D is generated using the baculovirus expression system BAC-TO-BAC (GIBCO BRL) based on *Autographa californica* nuclear polyhedrosis virus (AcNPV) infection of *Spodoptera frugiperda* insect cells (Sf9 cells).

25

cDNA encoding PDE is cloned into either the donor plasmid pFASTBAC1 or pFASTBAC-HT which contain a mini-Tn7 transposition element. The recombinant plasmid is transformed into DH10BAC competent cells which contain the parent bacmid bMON14272 (AcNPV infectious DNA) and a helper plasmid. The mini-Tn7 element on the pFASTBAC donor can transpose to the attTn7 attachment site on the bacmid thus introducing the PDE gene into the viral genome. Colonies containing recombinant bacmids are identified by disruption of the *lacZ* gene. The PDE/bacmid construct can then be isolated and infected into insect cells (Sf9 cells) resulting in the production of infectious recombinant baculovirus particles and expression of either

30



- 92 -

unfused recombinant enzyme (pFastbac1) or PDE4D-His fusion protein (pFastbacHT).

Cells are harvested and extracts prepared 24, 48 and 72 hours after transfection. Expression of PDE4D is confirmed by coomassie staining after sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) and western blotting onto a PVDF membrane of an unstained SDS-PAGE. The PDE-His fusion protein is detected due to the interaction between the Ni-NTA HRP conjugate and the His-tag which is fused to PDE4D.

#### Example 6: Production of PDE4D Specific Antibodies

Two approaches are utilized to raise antibodies to PDE4D, and each approach is useful for generating either polyclonal or monoclonal antibodies. In one approach, denatured protein from reverse phase HPLC separation is obtained in quantities up to 75 mg. This denatured protein is used to immunize mice or rabbits using standard protocols; about 100 µg are adequate for immunization of a mouse, while up to 1 mg might be used to immunize a rabbit. For identifying mouse hybridomas, the denatured protein is radioiodinated and used to screen potential murine B-cell hybridomas for those which produce antibody. This procedure requires only small quantities of protein, such that 20 mg is sufficient for labeling and screening of several thousand clones.

In the second approach, the amino acid sequence of an appropriate PDE4D domain, as deduced from translation of the cDNA, is analyzed to determine regions of high antigenicity. Oligopeptides comprising appropriate hydrophilic regions are synthesized and used in suitable immunization protocols to raise antibodies. The optimal amino acid sequences for immunization are usually at the C-terminus, the N-terminus and those intervening, hydrophilic regions of the polypeptide which are likely to be exposed to the external environment when the protein is in its natural conformation.

Typically, selected peptides, about 15 residues in length, are synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry and coupled to keyhole limpet hemocyanin (KLH; Sigma, St. Louis, MO) by reaction with M-maleimidobenzoyl-N-hydroxysuccinimide ester, MBS. If necessary, a cysteine is introduced at the N-terminus of the peptide to permit coupling to KLH. Rabbits are immunized with the peptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity by binding the peptide to plastic, blocking with 1% bovine serum albumin, reacting with antisera, washing and reacting with labeled (radioactive or fluorescent), affinity purified, specific goat anti-rabbit IgG.

Hybridomas are prepared and screened using standard techniques. Hybridomas of interest are detected by screening with labeled PDE4D to identify those fusions producing the monoclonal antibody with the desired specificity. In a typical protocol, wells of plates (FAST; Becton-Dickinson, Palo Alto, CA) are coated during incubation with affinity purified, specific rabbit anti-mouse (or suitable antispecies 1 g) antibodies at 10 mg/ml. The coated wells are blocked with 1% bovine serum albumin, (BSA), washed and incubated with supernatants from hybridomas. After washing the wells are incubated with labeled PDE4D at 1 mg/ml. Supernatants with specific antibodies bind more labeled PDE4D than is detectable in the background. Then clones producing specific antibodies are expanded and subjected to two cycles of cloning at limiting dilution. Cloned hybridomas are injected into pristane-treated mice to produce ascites, and monoclonal antibody is purified from mouse ascitic fluid by affinity chromatography on Protein A. Monoclonal antibodies with affinities of at least

$10^8 \text{ M}^{-1}$ , preferably  $10^9$  to  $10^{10} \text{ M}^{-1}$  or stronger, are typically made by standard procedures.

**Example 7: Diagnostic Test Using PDE4D Specific Antibodies**

Particular PDE4D antibodies are useful for investigating signal transduction and the diagnosis of infectious or hereditary conditions which are characterized by differences in the amount or distribution of PDE4D or downstream products of an active signaling cascade.

Diagnostic tests for PDE4D include methods utilizing antibody and a label to detect PDE4D in human body fluids, membranes, cells, tissues or extracts of such. The polypeptides and antibodies of the present invention are used with or without modification. Frequently, the polypeptides and antibodies are labeled by joining them, either covalently or noncovalently, with a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and have been reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent agents, chemiluminescent agents, chromogenic agents, magnetic particles and the like.

A variety of protocols for measuring soluble or membrane-bound PDE4D, using either polyclonal or monoclonal antibodies specific for the protein, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). A two-site monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on PDE4D is preferred, but a competitive binding assay may be employed.

**Example 8: Purification of Native PDE4D Using Specific Antibodies**

Native or recombinant PDE4D is purified by immunoaffinity chromatography using antibodies specific for PDE4D. In general, an immunoaffinity column is constructed by covalently coupling the anti-TRH antibody to an activated chromatographic resin.

- 95 -

Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated Sepharose (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

Such immunoaffinity columns are utilized in the purification of PDE4D by preparing a fraction from cells containing PDE4D in a soluble form. This preparation is derived by solubilization of whole cells or of a subcellular fraction obtained via differential centrifugation (with or without addition of detergent) or by other methods well known in the art. Alternatively, soluble PDE4D containing a signal sequence is secreted in useful quantity into the medium in which the cells are grown.

A soluble PDE4D-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PDE4D (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/protein binding (e.g., a buffer of pH 2-3 or a high concentration of a chaotrope such as urea or thiocyanate ion), and PDE4D is collected.

#### Example 9: Drug Screening

This invention is particularly useful for screening therapeutic compounds by using PDE4D or fragments thereof in any of a variety of drug screening techniques.

The following example provides a system for drug screening measuring the phosphodiesterase activity.

The phosphodiesterase activity of the crude extracts is measured and confirmed that the PDE cDNA encodes a phosphodiesterase which is able to hydrolyze cAMP or cGMP or both.

5 The recombinant PDE-His fusion protein can be purified from the crude lysate by metal-affinity chromatography using Ni-NTA agarose. This allows the specific retention of the recombinant material (since this is fused to the His-tag) whilst the endogenous insect proteins are washed off. The recombinant material is then eluted by competition with imidazol.

10 The phosphodiesterase activity of the recombinant protein is assayed using a commercially available SPA (scintillation proximity assay) kit (Amersham Pharmacia). The PDE enzyme hydrolyzes cyclic nucleotides, e.g. cAMP and cGMP to their linear counterparts. The SPA assay utilizes the tritiated cyclic nucleotides  
15 [ $^3\text{H}$ ]cAMP or [ $^3\text{H}$ ]cGMP, and is based upon the selective interaction of the tritiated non cyclic product with the SPA beads whereas the cyclic substrates are not effectively binding. Radiolabelled product bound to the scintillation beads generates light that can be analyzed in a scintillation counter.

## 20 Example 10: Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact, agonists, antagonists, or inhibitors. Any of these examples are used to fashion drugs  
25 which are more active or stable forms of the polypeptide or which enhance or interfere with the function of a polypeptide in vivo.

In one approach, the three-dimensional structure of a protein of interest, or of a protein-inhibitor complex, is determined by x-ray crystallography, by computer  
30 modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the polypeptide must be ascertained to elucidate the structure and to

- 97 -

determine active site(s) of the molecule. Less often, useful information regarding the structure of a polypeptide is gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design efficient inhibitors. Useful examples of rational drug design include molecules which have improved activity or stability or which act as inhibitors, agonists, or antagonists of native peptides.

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design is based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids is expected to be an analog of the original receptor. The anti-id is then used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides then act as the pharmacore.

By virtue of the present invention, sufficient amount of polypeptide are made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PDE4D amino acid sequence provided herein provides guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

#### **Example 11: Identification of Other Members of the Signal Transduction Complex**

Labeled PDE4D is useful as a reagent for the purification of molecules with which it interacts. In one embodiment of affinity purification, PDE4D is covalently coupled to a chromatography column. Cell-free extract derived from synovial cells or putative target cells is passed over the column, and molecules with appropriate affinity bind to PDE4D. PDE4D-complex is recovered from the column, and the

- 98 -

PDE4D-binding ligand disassociated and subjected to N-terminal protein sequencing. The amino acid sequence information is then used to identify the captured molecule or to design degenerate oligonucleotide probes for cloning the relevant gene from an appropriate cDNA library.

5

In an alternate method, antibodies are raised against PDE4D, specifically monoclonal antibodies. The monoclonal antibodies are screened to identify those which inhibit the binding of labeled PDE4D. These monoclonal antibodies are then used therapeutically.

10

#### **Example 12: Use and Administration of Antibodies, Inhibitors, or Antagonists**

15

Antibodies, inhibitors, or antagonists of PDE4D or other treatments and compounds that are limiters of signal transduction (LSTs), provide different effects when administered therapeutically. LSTs are formulated in a nontoxic, inert, pharmaceutically acceptable aqueous carrier medium preferably at a pH of about 5 to 8, more preferably 6 to 8, although pH may vary according to the characteristics of the antibody, inhibitor, or antagonist being formulated and the condition to be treated. Characteristics of LSTs include solubility of the molecule, its half-life and antigenicity/immunogenicity. These and other characteristics aid in defining an effective carrier. Native human proteins are preferred as LSTs, but organic or synthetic molecules resulting from drug screens are equally effective in particular situations.

20

25

LSTs are delivered by known routes of administration including but not limited to topical creams and gels; transmucosal spray and aerosol; transdermal patch and bandage; injectable, intravenous and lavage formulations; and orally administered liquids and pills particularly formulated to resist stomach acid and enzymes. The particular formulation, exact dosage, and route of administration is determined by the attending physician and varies according to each specific situation.

30

- 99 -

Such determinations are made by considering multiple variables such as the condition to be treated, the LST to be administered, and the pharmacokinetic profile of a particular LST. Additional factors which are taken into account include severity of the disease state, patient's age, weight, gender and diet, time and frequency of LST administration, possible combination with other drugs, reaction sensitivities, and tolerance/response to therapy. Long acting LST formulations might be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular LST.

Normal dosage amounts vary from 0.1 to  $10^5$   $\mu$ g, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. Those skilled in the art employ different formulations for different LSTs. Administration to cells such as nerve cells necessitates delivery in a manner different from that to other cells such as vascular endothelial cells.

It is contemplated that abnormal signal transduction, trauma, or diseases which trigger PDE4D activity are treatable with LSTs. These conditions or diseases are specifically diagnosed by the tests discussed above, and such testing should be performed in suspected cases of viral, bacterial or fungal infections, allergic responses, mechanical injury associated with trauma, hereditary diseases, lymphoma or carcinoma, or other conditions which activate the genes of lymphoid or neuronal tissues.

### **Example 13: Production of Non-human Transgenic Animals**

Animal model systems which elucidate the physiological and behavioral roles of the PDE4D are produced by creating nonhuman transgenic animals in which the activity of the PDE4D is either increased or decreased, or the amino acid sequence of the expressed PDE4D is altered, by a variety of techniques. Examples of these techniques include, but are not limited to: 1) Insertion of normal or mutant versions



- 100 -

of DNA encoding a PDE4D, by microinjection, electroporation, retroviral transfection or other means well known to those skilled in the art, into appropriately fertilized embryos in order to produce a transgenic animal or 2) homologous recombination of mutant or normal, human or animal versions of these genes with the native gene locus in transgenic animals to alter the regulation of expression or the structure of these PDE4D sequences. The technique of homologous recombination is well known in the art. It replaces the native gene with the inserted gene and hence is useful for producing an animal that cannot express native PDE4Ds but does express, for example, an inserted mutant PDE4D, which has replaced the native PDE4D in the animal's genome by recombination, resulting in underexpression of the transporter. Microinjection adds genes to the genome, but does not remove them, and the technique is useful for producing an animal which expresses its own and added PDE4D, resulting in overexpression of the PDE4D.

One means available for producing a transgenic animal, with a mouse as an example, is as follows: Female mice are mated, and the resulting fertilized eggs are dissected out of their oviducts. The eggs are stored in an appropriate medium such as cesiumchloride M2 medium. DNA or cDNA encoding PDE4D is purified from a vector by methods well known to the one skilled in the art. Inducible promoters may be fused with the coding region of the DNA to provide an experimental means to regulate expression of the transgene. Alternatively or in addition, tissue specific regulatory elements may be fused with the coding region to permit tissue-specific expression of the transgene. The DNA, in an appropriately buffered solution, is put into a microinjection needle (which may be made from capillary tubing using a piper puller) and the egg to be injected is put in a depression slide. The needle is inserted into the pronucleus of the egg, and the DNA solution is injected. The injected egg is then transferred into the oviduct of a pseudopregnant mouse which is a mouse stimulated by the appropriate hormones in order to maintain false pregnancy, where it proceeds to the uterus, implants, and develops to term. As noted above, microinjection is not the only method for inserting DNA into the egg but is used here only for exemplary purposes.

References

- U.S. 4,522,811
- 5 U.S. 5,283,317
- U.S. 5,565,332
- U.S. 5,977,305
- WO 84/03564
- WO 92/01810
- 10 WO 93/03151
- WO 94/13804
- WO 01/04297
- WO 01/60860
- WO 02/28999
- 15 Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ;  
Nucleic Acids Res 1997 Sep 1; 25(17): 3389-402
- Appa Rao et al., 1997, Protein Expr Purif Nov, 11(2): 201-8
- Baecker PA et al., Gene 1994 Jan 28;138(1-2):253-6
- Bang, Y.J. et al. (1994)Proc. Natl. Acad. Sci. USA 91:5330-5334
- 20 Bardelle, C. et al. (1999) Anal. Biochem. 275: 148-155
- Barnes, 2000, Chest, 117:10S14S
- Beavo, Physiol. Rev . 75:725-748 (1995);
- Botstein et al., 1980, Am J Hum Genet. 32: 314-31
- Charbonneau, et al., Proc. Natl. Acad. Sci . ( USA ) 83:9308-9312 (1986)
- 25 Colbere-Garapin *et al.*, 1981, *J. Mol. Biol.* 150, 1-14
- Conti, et al., Physiol. Rev . 75:723-748 (1995)].
- Conti et al. (1995) Endocrine Rev. 16:370-389
- Engelhard *et al.*, 1994, *Proc. Nat. Acad. Sci.* 91, 3224-3227
- Francis, et al., J. Biol. Chem . 269:22477-22480 (1994).
- 30 Fujishige, K. et al. (1999) J. Biol. Chem. 274:18438-18445
- Gergen and Weiss , 1992, Am Rev Respir Dis 146:823-824

- 102 -

- Gibson et al., 1996, *Genome Research* 6: 995-1001
- Han, P. et al. (1997) *J. Biol. Chem.* 272:16152-16157
- Haseloff *et al.*, 1988, *Nature* 334, 585-591
- Harbinson, et al., *Eur. Respir. J.* 10:1008-1014 (1997)
- 5 Heid et al., 1996, *Genome Research* 6: 986-994
- Hetman JM, Robas N, Baxendale R, Fidock M, Phillips SC, Soderling SH, Beavo JA  
*Proc. Natl. Acad. Sci. U S A* 2000 Nov 7;97(23):12891-5
- Hansen, R.S., and Beavo, J.A., *PNAS USA* 1982;79: 2788-92
- Holland et al., 1991, *PNAS* 88: 7276-7280
- 10 Iwabuchi *et al.*, 1993, *Oncogene* 8, 1693-1696
- Jeffreys et al., 1985, *Nature* 316: 76-9
- Johnson *et al.*, 1989, *Endoc. Rev.* 10, 317-331
- Kakkar, R. et al. (1999) *Cell Mol Life Sci* 55:1164-1186
- Kellogg et al., 1990, *Anal. Biochem.* 189:202-208
- 15 Lam, 1997, *Anticancer Drug Res.* 12(3):145-67
- Le Jeune IR et al., *J Biol Chem* 2002 Sep 27;277(39):35980-9
- Livak et al., 1995, *PCR Methods and Applications* 357-362
- Logan, Shenk, 1984, *Proc. Natl. Acad. Sci.* 81, 3655-3659
- Loughney and Ferguson, in *Phosphodiesterase Inhibitors*, Schudt, et al. (Eds.),  
Academic Press: New York, N.Y. (1996) pp. 1- 19
- 20 Loughney, K. et al (1999) *Gene* 234:109-117
- Lowy *et al.*, 1980, *Cell* 22, 817-23
- Maddox *et al.*, 1983, *J. Exp. Med.* 158, 1211-1216
- McAllister-Lucas, L.M. (1995) *J. Biol. Chem.* 270:30671-30679
- 25 McConnell *et al.*, 1992, *Science* 257, 1906-1912
- Manganiello, et al., *Arch. Biochem. Acta* 322:1-13 (1995);
- Nemoz G et al., *FEBS Lett* 1996 Apr 8;384(1):97-102
- Nicholls *et al.*, 1993, *J. Immunol. Meth.* 165, 81-91
- Perry, M.J. and G.A. Higgs (1998) *Curr. Opin. Chem. Biol.* 2:472-481
- 30 Piatak et al., 1993, *BioTechniques* 14:70-81
- Piatak et al., 1993, *Science* 259:1749-1754

- 103 -

- Porath *et al.*, 1992, *Prot. Exp. Purif.* 3, 263-281
- Roberge *et al.*, 1995, *Science* 269, 202-204
- Sadhu, K. *et al.* (1999) *J. Histochem Cytochem* 47:895-906
- Scott and Smith (1990) *Science* 249:386-390
- 5 Sjolander, Urbaniczky, 1991, *Anal. Chem.* 63, 2338-2345
- Soderling, S.H. *et al.* (1999) *Proc. Natl. Acad. Sci. USA* 96:7071-7076
- Szabo *et al.*, 1995, *Curr. Opin. Struct. Biol.* 5, 699-705
- Terrett, N. *et al.* (1996) *Bioorg. Med. Chem. Lett.* 6:1819-1824
- Thomas, 1980, *Proc. Nat. Acad. Sci.*, 77:5201-5205
- 10 Turko, I.V. *et al.* (1998) *Biochemistry* 37:4200-4205
- Uhlmann *et al.*, 1987, *Tetrahedron. Lett.* 215, 3539-3542
- Weber *et al.*, 1990, *Genomics* 7: 524-30
- Wigler *et al.*, 1977, *Cell* 11, 223-32
- Wigler *et al.*, 1980, *Proc. Natl. Acad. Sci.* 77, 3567-70

15

Claims

1. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising the steps of
- 5
- i) contacting a test compound with a PDE4D polypeptide,
- 10
- ii) detect binding of said test compound to said PDE4D polypeptide.
2. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising the steps of
- 15
- i) determining the activity of a PDE4D polypeptide at a certain concentration of a test compound or in the absence of said test compound,
- 20
- ii) determining the activity of said polypeptide at a different concentration of said test compound.
- 25
3. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising the steps of
- 30

- 105 -

- i) determining the activity of a PDE4D polypeptide at a certain concentration of a test compound,
  - ii) determining the activity of a PDE4D polypeptide at the presence of a compound known to be a regulator of a PDE4D polypeptide.
- 5
- 4. The method of any of claims 1 to 3, wherein the step of contacting is in or at the surface of a cell.
  - 5. The method of any of claims 1 to 3, wherein the cell is in vitro.
  - 6. The method of any of claims 1 to 3, wherein the step of contacting is in a cell-free system.
  - 7. The method of any of claims 1 to 3, wherein the polypeptide is coupled to a detectable label.
  - 8. The method of any of claims 1 to 3, wherein the compound is coupled to a detectable label.
  - 9. The method of any of claims 1 to 3, wherein the test compound displaces a ligand which is first bound to the polypeptide.
  - 10. The method of any of claims 1 to 3, wherein the polypeptide is attached to a solid support.
  - 11. The method of any of claims 1 to 3, wherein the compound is attached to a solid support.
  - 12. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of disorders of the
- 10
- 15
- 20
- 25
- 30

- 106 -

peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising the steps of

- 5           i)       contacting a test compound with a PDE4D polynucleotide,
- ii)       detect binding of said test compound to said PDE4D polynucleotide.

13.       The method of claim 12 wherein the nucleic acid molecule is RNA.

10

14.       The method of claim 12 wherein the contacting step is in or at the surface of a cell.

15.       The method of claim 12 wherein the contacting step is in a cell-free system.

15

16.       The method of claim 12 wherein polynucleotide is coupled to a detectable label.

17.       The method of claim 12 wherein the test compound is coupled to a detectable label.

20

18.       A method of diagnosing a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising the steps of

25

i)       determining the amount of a PDE4D polynucleotide in a sample taken from said mammal,

30

ii)       determining the amount of PDE4D polynucleotide in healthy and/or diseased mammals.

- 107 -

19. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising a therapeutic agent which binds to a PDE4D polypeptide.
20. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising a therapeutic agent which regulates the activity of a PDE4D polypeptide.
21. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising a therapeutic agent which regulates the activity of a PDE4D polypeptide, wherein said therapeutic agent is
- i) a small molecule,
  - ii) an RNA molecule,
  - iii) an antisense oligonucleotide,
  - iv) a polypeptide,
  - v) an antibody, or
  - vi) a ribozyme.
22. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising a PDE4D polynucleotide.



23. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising a PDE4D polypeptide.
24. Use of regulators of a PDE4D for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal.
25. Method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal comprising the steps of
- i) identifying a regulator of PDE4D,
  - ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders in a mammal; and
  - iii) combining of said regulator with an acceptable pharmaceutical carrier.
26. Use of a regulator of PDE4D for the regulation of PDE4D activity in a mammal having a disease comprised in a group of diseases consisting of disorders of the peripheral and central nervous system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders.

- 1/5 -

Fig. 1

SEQ ID NO: 1

GAATTCCCTGGATACAGCCTTTTTATGACTTTTACTTCCTTTATANNC  
AAATTCCAACGTCTTCTCATTCCTCCACCAGGSCTGTGCCAACCTGGG  
CCCAACCCAAGGSCCTCACTAAACCATCCAATCAGTAGGAGCCATAGA  
CTACTTTATTTAGCCAAAGCAAAAATGAGTCAACTGAATTCTGTTTTT  
CCATTTACTTCTGTCTGTTTTTTCCTTCCTCTTGCCACCCTCAGTGCCA  
CAAGAGGGGACCCCTCTCGGTAGCCCTGAGGCTCTGGCGCCTTCAAGT  
GAGAAGCTAAGCACCAGCCTCTGCTGGGCTGCAGAAGCGGCGGCGGCG  
GCAGCAGCAGCAGCAGCATCAGGAAGGCTCTCGGGCCAGCGCGGTGAA  
CCCGGGCTGGGCAGCAGGTCGCGGAGCCGCGAGCCAGGATGGAGGCAG  
AGGGCAGCAGCGCGCCGGCCCCGGGCGGGCAGCGGAGAGGGCAGCGACA  
GCGCCGGCGGGGCCACGCTCAAAGCCCCAAGCATCTCTGGAGGCACG  
AGCAGCACCACCAGTACCCGCTCCGGCAGCCCCAGTTCCGCCTCCTGC  
ATCCCCATCACCACCTGCCCCCGCCGCGCCACCCTCGCCCCAGCCCC  
AGCCCCAGTGTCCGCTACAGCCGCGCGCGCGCCCCCCCCCTGCCGCCGC  
CCCCGCCGCGCGCCGGGGCTGCCCGCGGCGGCTACGCCTCGAGCGGGG  
CCACCGGCCGCGTCCGGCATCGCGGCTACTCGGACACCGAGCGCTACC  
TGTA CTGTGCGGCCATGGACCGCACCTCCTACGCGGTGGAGACCGGCC  
ACCGGCCCGCCTGAAGAAATCCAGGATGTCCTGGCCCTCCTCGTTCC  
AGGGACTCAGGCGTTTTGATGTGGACAATGGCACATCTGCGGGACGGA  
GTCCCTTGATCCCATGACCAGCCCAGGATCCGGGCTAATTCTCCAAG  
CAAATTTTGTCCACAGTCAACGACGGGAGTCCTTCCTGTATCGATCCG  
ACAGCGATTATGACCTCTCTCAAAGTCTATGTCCCGGAACTCCTCCA  
TTGCCAGTGATATACAGGAGATGACTTGATTGTGACTCCATTTGCTC  
AGGTCTTGGCCAGTCTGCGAACTGTACGAAACAACTTTGCTGCATTAA  
CTAATTTGCAAGATCGAGCACCTAGCAAAAGATCACCCATGTGCAACC  
AACCATCCATCAACAAAGCCACCATAACAGAGGAGGCCTACCAGAAAC  
TGGCCAGCGAGACCCTGGAGGAGCTGGACTGGTGTCTGGACCAGCTAG

- 2/5 -

Fig. 1 (continued)

AGACCCTACAGACCAGGCACTCCGTCAGTGAGATGGCCTCCAACAAGT  
TTAAAAGGATGCTTAATCGGGAGCTCACCCATCTCTCTGAAATGAGTC  
GGTCTGGAAATCAAGTGTGAGAGTTTATATCAAACACATTCTTAGATA  
AGCAACATGAAGTGGAAATTCCTTCTCCAACCTCAGAAGGAAAAGGAGA  
AAAAGAAAAGACCAATGTCTCAGATCAGTGGAGTCAAGAAATTGATGC  
ACAGCTCTAGTCTGACTAATTCAAGTATCCCAAGGTTTGGAGTTAAAA  
CTGAACAAGAAGATGTCCTTGCCAAGGAAC TAGAAGATGTGAACAAAT  
GGGGTCTTCATGTTTTTCAGAATAGCAGAGTTGTCTGGTAACCGGCCCT  
TGACTGTTATCATGCACACCATTTTTTCAGGAACGGGATTTATTAAAAA  
CATTTAAAATTCCAGTAGATACTTTAATTACATATCTTATGACTCTCG  
AAGACCATTACCATGCTGATGTGGCCTATCACAACAATATCCATGCTG  
CAGATGTTGTCCAGTCTACTCATGTGCTATTATCTACACCTGCTTTGG  
AGGCTGTGTTTACAGATTTGGAGATTCTTGCAGCAATTTTTTGCCAGTG  
CAATACATGATGTAGATCATCCTGGTGTGTCCAATCAATTTCTGATCA  
ATACAAACTCTGAACTTGCCTTGATGTACAATGATTCCTCAGTCTTAG  
AGAACCATCATTTGGCTGTGGGCTTTAAATTGCTTCAGGAAGAAAAC  
GTGACATTTTCCAGAATTTGACCAAAAAACAAAGACAATCTTTAAGGA  
AAATGGTCATTGACATCGTACTTGCAACAGATATGTCAAAACACATGA  
ATCTACTGGCTGATTTGAAGACTATGGTTGAAACTAAGAAAGTGACAA  
GCTCTGGAGTTCTTCTTCTTGATAATTATTCCGATAGGATTCAGGTTC  
TTCAGAATATGGTGCACTGTGCAGATCTGAGCAACCCAACAAAGCCTC  
TCCAGCTGTACCGCCAGTGGACGGACCGGATAATGGAGGAGTTCTTCC  
GCCAAGGAGACCGAGAGAGGGAACGTGGCATGGAGATAAGCCCCATGT  
GTGACAAGCACAATGCTTCCGTGGAAAAATCACAGGTGGGCTTCATAG  
ACTATATTGTTTCATCCCCTCTGGGAGACATGGGCAGACCTCGTCCACC  
CTGACGCCCAGGATATTTTGGACACTTTGGAGGACAATCGTGAATGGT  
ACCAGAGCACAATCCCTCAGAGCCCCTCTCCTGCACCTGATGACCCAG

- 3/5 -

Fig. 1 (continued)

AGGAGGGCCGGCAGGGTCAAACCTGAGAAATTCAGTTTGAACCTAACTT  
TAGAGGAAGATGGTGAGTCAGACACGGAAAAGGACAGTGGCAGTCAAG  
TGGAAGAAGACACTAGCTGCAGTGACTCCAAGACTCTTTGTACTCAAG  
ACTCAGAGTCTACTGAAATTTCCCCTTGATGAACAGGTTGAAGAGGAGG  
CAGTAGGGGAAGAAGAGGAAAGCCAGCCTGAAGCCTGTGTCATAGATG  
ATCGTTCTCCTGACACGTAACAGTGCAAAAACCTTTCATGCCTTTTTTT  
TTTTTAAGTAGAAAAATTGTTTCCAAAGTGTCATGTCACATGCCACAAC  
CACGGTCACACCTCACTGTCATCTGCCAGGACGTTTGTGTAACAAAAC  
TGACCTTGACTACTCAGTCCAGCGCTCAGGAATATCGTAACCAGTTTT  
TTCACCTCCATGTTTCATCCGAGCAAGGTGGACATCTTCACGAACAGCG  
TTTTTAACAAGATTTTCAGCTTGGTAGAGCTGACAAAGCAGATAAAATC  
TACTCCAAATTATTTTCAAGAGAGTGTGACTCATCAGGCAGCCCCAAA  
GTTTATTGGACTTGGGGTTTCTATTCTTTTTTATTTGTTTGCAATATT  
TTCAGAAGAAAGGCATTGCACAGAGTGAACCTTAATGGACGAAGCAACA  
AATATGTCAAGAACAGGACATAGCACGAATCTGTTACCAGTAGGAGGA  
GGATGAGCCACAGAAATTGCATAATTTTCTAATTTCAAGTCTTCCTGA  
TACATGACTGAATAGTGTGGTTCAGTGAGCTGCACTGACCTCTACATT  
TTGTATGATATGTAAACAGATTTTTTGTAGAGCTTACTTTTATTATT  
AAATGTATTGAGGTATTATATTTAAAAAAACTATGTTTCAGAACTTCA  
TCTGCCACTGGTTATTTTTTTTCTAAGGAGTAACTTGCAAGTTTTTCAGT  
ACAAATCTGTGCTACACTGGATAAAAATCTAATTTATGAATTTTACTT  
GCACCTTATAGTTCATAGCAATTAACCTGATTTGTAGTGATTCATTGTT  
TGTTTTATATACCAATGACTTCCATATTTTAAAAGAGAAAAACAACTT  
TATGTTGCAGGAAACCCTTTTTGTAAAGTCTTTATTATTTACTTTGCAT  
TTTGTTTCACTCTTTCCAGATAAGCAGAGTTGCTCTTCACCAGTGTTT  
TTCTTCATGTGCAAAGTGACTATTTGTTCTATAATAC

- 4/5 -

Fig. 2

SEQ ID NO: 2

MEAEGSSAPARAGSGEGSDSAGGATLKAPKHLWRHEQHHQYPLRQPQF  
RLHHPHHHLPPPPPPSPQPQPQCPLQPPPPPLPPPPPPGAARGRYA  
SSGATGRVRHRGYSDTERYLYCRAMDRTSYAVETGHRPGLKKSRRMSWP  
SSFQGLRRFDVDNGTSAGRSPLDPMTSPGSGLILQANFVHSQRRESFL  
YRSDSDYDLSPKSMRNSSIASDIHGDDLIVTPFAQVLASLRTVRNNF  
AALTNLQDRAPSKRSPMCNQPSINKATITTEEAYQKLASETLEELDWC  
DQLETLQTRHSVSEMASNKFKRMLNRELTHLSEMSRSGNQVSEFISNT  
FLDKQHEVEIPSPTQKEKEKKRPMSSQISGVKKLMHSSSLTNSSIPRF  
GVKTEQEDVLAKELEDVNKWGLHVFRIAELSGNRPLTVIMHTIFQERD  
LLKTFKIPVDTLITYLMTLEDHYHADVAYHNNIHAADVQSTHVLLST  
PALEAVFTDLEILAAIFASAIHDVDHPGVSNOFLINTNSELALMYNDS  
SVLENHHLAVGFKLLQEENCDFQNLTKKQRQSLRKMVIDIVLATDMS  
KHMNLLADLKTMTVETKKVTSSGVLLLDNYSDRIQVLQNMVHCADLSNP  
TKPLQLYRQWTDRIEIEFFRQGDREMERGMEISPMCDKHNASVEKSQV  
GFIDYIVHPLWETWADLVHPDAQDILDTLEDNREWYQSTIPQSPSPAP  
DDPEEGRQGQTEKFQFELTLEEDGESDTEKDSGSQVEEDTSCSDSKTL  
CTQDSESTEIPLDEQVEEEEAVGEEESQPEACVIDDRSPDT

Fig. 3

SEQ ID NO: 3

5' -GAATATGGTGCACTGTGCAGATCT-3'

Fig. 4

SEQ ID NO: 4

5' -TCCATTATCCGGTCCGTCC-3'

- 5/5 -

Fig. 5

SEQ ID NO: 5

5' - CAAAGCCTCTCCAGCTGTACCGCCA - 3'

- 1 -

## SEQUENCE LISTING

&lt;110&gt; Bayer, BHC

<120> Diagnostics and Therapeutics for Diseases Associated with Human  
Phosphodiesterase 4D (PDE4D)

&lt;130&gt; Le A 36 430

&lt;160&gt; 5

&lt;170&gt; PatentIn version 3.1

&lt;210&gt; 1

&lt;211&gt; 3829

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (46)..(46)

&lt;223&gt; n=a,t,g,c

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (47)..(47)

&lt;223&gt; n=a,t,g,c

- 2 -

<400> 1  
gaattccctg gatacagcct ttttatgact tttacttcct ttatanncaa attccaacgt 60  
cttctcattc ctccaccagg sctgtgccaa cctgggcca acccaaggsc ctactaaac 120  
catccaatca gtaggagcca tagactactt tatttagcca aagcaaaaat gagtcaactg 180  
aattctgttt ttccatttac ttctgtctgt ttttcttcc tottgccacc ctcatgcca 240  
caagagggga cccctctcgg tagccctgag gctctggcgc cttcaagtga gaagctaagc 300  
accagcctct gctgggctgc agaagcggcg gcggcgagc cagcagcagc agcatcagga 360  
aggctctcgg gccagcggg tgaacccggg ctgggcagca ggtcgggag ccgcgagcca 420  
ggatggaggc agagggcagc agcgcggcgg cccgggcggg cagcggagag ggcagcgaca 480  
gcgcggcggg ggccacgctc aaagcccca agcatctctg gaggcacgag cagcaccacc 540  
agtaccgct ccggcagccc cagtccgcc tcctgcatcc ccatcaccac ctgccccgc 600  
cgccgccacc ctgcgccag cccagcccc agtgtccgct acagccgccg ccgcggcccc 660  
ccctgccgcc gccccgcg ccgcccgggg ctgcccggcg ccgctacgcc tcgagcgggg 720  
ccaccggcgg cgtccggcat cggggtact cggacaccga gcgctacctg tactgtcgcg 780  
ccatggaccg cacctcctac gcggtggaga ccggccaccg gcccggcctg aagaaatcca 840  
ggatgtcctg gccctcctcg ttccaggac tcaggcggtt tgatgtggac aatggcacat 900  
ctgcgggacg gagtcccttg gatcccatga ccagcccagg atccgggcta attctccaag 960  
caaattttgt ccacagtcaa cgaaggaggc ccttctgta tcgatccgac agcgattatg 1020  
acctctctcc aaagtctatg tcccgaact cctccattgc cagtgatata cacggagatg 1080  
acttgattgt gactccattt gctcaggctt tggccagtct gcgaactgta cgaaacaact 1140  
ttgctgcatt aactaatttg caagatcgag cacctagcaa aagatcacc atgtgcaacc 1200  
aaccatccat caacaaagcc accataacag aggaggccta ccagaaactg gccagcgaga 1260  
ccctggagga gctggactgg tgtctggacc agctagagac cctacagacc aggcactccg 1320  
tcagtggagat ggctccaac aagtttaaaa ggatgcttaa tcgggagctc acccatctct 1380



- 3 -

ctgaaatgag tcggtctgga aatcaagtgt cagagtttat atcaaacaca ttcttagata 1440

agcaacatga agtggaaatt ctttctccaa ctccagaagga aaaggagaaa aagaaaagac 1500

caatgtctca gatcagtgga gtcaagaaat tgatgcacag ctctagtctg actaattcaa 1560

gtatcccaag gtttggagtt aaaactgaac aagaagatgt ccttgccaag gaactagaag 1620

atgtgaacaa atgggggtctt catgttttca gaatagcaga gttgtctggt aaccggccct 1680

tgactgttat catgcacacc atttttcagg aacgggattt attaaaaaca tttaaaattc 1740

cagtagatac ttttaattaca tatcttatga ctctcgaaga ccattaccat gctgatgtgg 1800

cctatcacia caatatccat gctgcagatg ttgtccagtc tactcatgtg ctattatcta 1860

cacctgcttt ggaggctgtg tttacagatt tggagattct tgcagcaatt tttgccagtg 1920

caatacatga tgtagatcat cctggtgtgt ccaatcaatt tctgatcaat acaaactctg 1980

aacttgctt gatgtacaat gattcctcag tcttagagaa ccatcatttg gctgtgggct 2040

ttaaattgct tcaggaagaa aactgtgaca ttttccagaa ttgaccaa aaacaaagac 2100

aatctttaag gaaaatggc attgacatcg tacttgcaac agatatgtca aaacacatga 2160

atctactggc tgatttgaag actatggttg aaactaagaa agtgacaagc tctggagttc 2220

ttcttcttga taattattcc gataggattc aggttcttca gaatatggtg cactgtgcag 2280

atctgagcaa cccaacaaag cctctccagc tgtaccgcca gtggacggac cggataatgg 2340

aggagtctt cgcgaagga gaccgagaga gggacgtgg catggagata agcccatgt 2400

gtgacaagca caatgcttcc gtggaaaaat cacagggtgg cttcatagac tatattgttc 2460

atccccctg ggagacatgg gcagacctcg tccacctga cggccaggat attttgga 2520

ctttggagga caatcgtgaa tgggtaccaga gcacaatccc tcagagcccc tctcctgcac 2580

ctgatgaccc agaggagggc cggcagggc aaactgagaa attccagttt gaactaactt 2640

tagaggaaga tggtagtca gacacggaaa aggacagtgg cagtcaagtg gaagaagaca 2700

ctagctgcag tgactccaag actctttgta ctcaagactc agagtctact gaaattcccc 2760

- 4 -

ttgatgaaca ggttgaagag gaggcagtag gggagaaga ggaaagccag cctgaagcct 2820  
 gtgtcataga tgatcggtct cctgacacgt aacagtgcaa aaactttcat gccttttttt 2880  
 tttttaagta gaaaaattgt ttccaaagtg catgtcacat gccacaacca cggtcacacc 2940  
 tcaactgtcat ctgccaggac gtttgttgaa caaaactgac cttgactact cagtccagcg 3000  
 ctgaggaata tcgtaaccag ttttttcacc tccatgttca tccgagcaag gtggacatct 3060  
 tcacgaacag cgtttttaac aagatttcag cttggtagag ctgacaaagc agataaaatc 3120  
 tactccaaat tattttcaag agagtgtgac tcatcaggca gcccaaaagt ttattggact 3180  
 tgggggtttct attccttttt atttgtttgc aatattttca gaagaaaggc attgcacaga 3240  
 gtgaacttaa tggacgaagc aacaaatatg tcaagaacag gacatagcac gaatctgtta 3300  
 ccagtaggag gaggatgagc cacagaaatt gcataatttt ctaatttcaa gtcttcctga 3360  
 tacatgactg aatagtgtgg ttcagtgagc tgcactgacc tctacatttt gtatgatatg 3420  
 taaaacagat tttttgtaga gcttactttt attattaaat gtattgaggt attatattta 3480  
 aaaaaaacta tgttcagaac ttcactgtcc actggttatt tttttctaag gagtaacttg 3540  
 caagttttca gtacaaatct gtgctacact ggataaaaat ctaatttatg aattttactt 3600  
 gcaccttata gttcatagca attaactgat ttgtagtgat tcattgtttg ttttatatac 3660  
 caatgacttc catattttta aagagaaaaa caactttatg ttgcaggaaa ccctttttgt 3720  
 aagtctttat tatttacttt gcattttgtt tcactctttc cagataagca gagttgctct 3780  
 tcaccagtgt ttttcttcat gtgcaaagtg actatttggt ctataatac 3829

&lt;210&gt; 2

&lt;211&gt; 809

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

- 5 -

&lt;400&gt; 2

Met Glu Ala Glu Gly Ser Ser Ala Pro Ala Arg Ala Gly Ser Gly Glu  
1 5 10 15

Gly Ser Asp Ser Ala Gly Gly Ala Thr Leu Lys Ala Pro Lys His Leu  
20 25 30

Trp Arg His Glu Gln His His Gln Tyr Pro Leu Arg Gln Pro Gln Phe  
35 40 45

Arg Leu Leu His Pro His His His Leu Pro Pro Pro Pro Pro Pro Ser  
50 55 60

Pro Gln Pro Gln Pro Gln Cys Pro Leu Gln Pro Pro Pro Pro Pro Pro  
65 70 75 80

Leu Pro Pro Pro Pro Pro Pro Pro Gly Ala Ala Arg Gly Arg Tyr Ala  
85 90 95

Ser Ser Gly Ala Thr Gly Arg Val Arg His Arg Gly Tyr Ser Asp Thr  
100 105 110

Glu Arg Tyr Leu Tyr Cys Arg Ala Met Asp Arg Thr Ser Tyr Ala Val  
115 120 125

Glu Thr Gly His Arg Pro Gly Leu Lys Lys Ser Arg Met Ser Trp Pro  
130 135 140

Ser Ser Phe Gln Gly Leu Arg Arg Phe Asp Val Asp Asn Gly Thr Ser  
145 150 155 160

Ala Gly Arg Ser Pro Leu Asp Pro Met Thr Ser Pro Gly Ser Gly Leu  
165 170 175

Ile Leu Gln Ala Asn Phe Val His Ser Gln Arg Arg Glu Ser Phe Leu  
180 185 190

Tyr Arg Ser Asp Ser Asp Tyr Asp Leu Ser Pro Lys Ser Met Ser Arg  
195 200 205

Asn Ser Ser Ile Ala Ser Asp Ile His Gly Asp Asp Leu Ile Val Thr  
210 215 220

- 6 -

Pro Phe Ala Gln Val Leu Ala Ser Leu Arg Thr Val Arg Asn Asn Phe  
 225 230 235 240

Ala Ala Leu Thr Asn Leu Gln Asp Arg Ala Pro Ser Lys Arg Ser Pro  
 245 250 255

Met Cys Asn Gln Pro Ser Ile Asn Lys Ala Thr Ile Thr Glu Glu Ala  
 260 265 270

Tyr Gln Lys Leu Ala Ser Glu Thr Leu Glu Glu Leu Asp Trp Cys Leu  
 275 280 285

Asp Gln Leu Glu Thr Leu Gln Thr Arg His Ser Val Ser Glu Met Ala  
 290 295 300

Ser Asn Lys Phe Lys Arg Met Leu Asn Arg Glu Leu Thr His Leu Ser  
 305 310 315 320

Glu Met Ser Arg Ser Gly Asn Gln Val Ser Glu Phe Ile Ser Asn Thr  
 325 330 335

Phe Leu Asp Lys Gln His Glu Val Glu Ile Pro Ser Pro Thr Gln Lys  
 340 345 350

Glu Lys Glu Lys Lys Lys Arg Pro Met Ser Gln Ile Ser Gly Val Lys  
 355 360 365

Lys Leu Met His Ser Ser Ser Leu Thr Asn Ser Ser Ile Pro Arg Phe  
 370 375 380

Gly Val Lys Thr Glu Gln Glu Asp Val Leu Ala Lys Glu Leu Glu Asp  
 385 390 395 400

Val Asn Lys Trp Gly Leu His Val Phe Arg Ile Ala Glu Leu Ser Gly  
 405 410 415

Asn Arg Pro Leu Thr Val Ile Met His Thr Ile Phe Gln Glu Arg Asp  
 420 425 430

Leu Leu Lys Thr Phe Lys Ile Pro Val Asp Thr Leu Ile Thr Tyr Leu  
 435 440 445

Met Thr Leu Glu Asp His Tyr His Ala Asp Val Ala Tyr His Asn Asn  
 450 455 460

- 7 -

Ile His Ala Ala Asp Val Val Gln Ser Thr His Val Leu Leu Ser Thr  
 465 470 475 480

Pro Ala Leu Glu Ala Val Phe Thr Asp Leu Glu Ile Leu Ala Ala Ile  
 485 490 495

Phe Ala Ser Ala Ile His Asp Val Asp His Pro Gly Val Ser Asn Gln  
 500 505 510

Phe Leu Ile Asn Thr Asn Ser Glu Leu Ala Leu Met Tyr Asn Asp Ser  
 515 520 525

Ser Val Leu Glu Asn His His Leu Ala Val Gly Phe Lys Leu Leu Gln  
 530 535 540

Glu Glu Asn Cys Asp Ile Phe Gln Asn Leu Thr Lys Lys Gln Arg Gln  
 545 550 555 560

Ser Leu Arg Lys Met Val Ile Asp Ile Val Leu Ala Thr Asp Met Ser  
 565 570 575

Lys His Met Asn Leu Leu Ala Asp Leu Lys Thr Met Val Glu Thr Lys  
 580 585 590

Lys Val Thr Ser Ser Gly Val Leu Leu Leu Asp Asn Tyr Ser Asp Arg  
 595 600 605

Ile Gln Val Leu Gln Asn Met Val His Cys Ala Asp Leu Ser Asn Pro  
 610 615 620

Thr Lys Pro Leu Gln Leu Tyr Arg Gln Trp Thr Asp Arg Ile Met Glu  
 625 630 635 640

Glu Phe Phe Arg Gln Gly Asp Arg Glu Arg Glu Arg Gly Met Glu Ile  
 645 650 655

Ser Pro Met Cys Asp Lys His Asn Ala Ser Val Glu Lys Ser Gln Val  
 660 665 670

Gly Phe Ile Asp Tyr Ile Val His Pro Leu Trp Glu Thr Trp Ala Asp  
 675 680 685

Leu Val His Pro Asp Ala Gln Asp Ile Leu Asp Thr Leu Glu Asp Asn  
 690 695 700

- 8 -

Arg Glu Trp Tyr Gln Ser Thr Ile Pro Gln Ser Pro Ser Pro Ala Pro  
 705 710 715 720

Asp Asp Pro Glu Glu Gly Arg Gln Gly Gln Thr Glu Lys Phe Gln Phe  
 725 730 735

Glu Leu Thr Leu Glu Glu Asp Gly Glu Ser Asp Thr Glu Lys Asp Ser  
 740 745 750

Gly Ser Gln Val Glu Glu Asp Thr Ser Cys Ser Asp Ser Lys Thr Leu  
 755 760 765

Cys Thr Gln Asp Ser Glu Ser Thr Glu Ile Pro Leu Asp Glu Gln Val  
 770 775 780

Glu Glu Glu Ala Val Gly Glu Glu Glu Glu Ser Gln Pro Glu Ala Cys  
 785 790 795 800

Val Ile Asp Asp Arg Ser Pro Asp Thr  
 805

<210> 3

<211> 24

<212> DNA

<213> Homo sapiens

<400> 3

gaatatggtg cactgtgcag atct

24

<210> 4

<211> 19

<212> DNA

<213> Homo sapiens

<400> 4

tccattatcc ggtccgtcc

19

- 9 -

&lt;210&gt; 5

&lt;211&gt; 25

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 5

caaagcctct ccagctgtac cgcca

25

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
21 May 2004 (21.05.2004)

PCT

(10) International Publication Number  
**WO 2004/042389 A3**

(51) International Patent Classification<sup>7</sup>: C12Q 1/44

(21) International Application Number:  
PCT/EP2003/011877

(22) International Filing Date: 25 October 2003 (25.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
02024993.4 8 November 2002 (08.11.2002) EP

(71) Applicant (for all designated States except US): BAYER  
HEALTHCARE AG [DE/DE]; 51368 Leverkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GOLZ, Stefan  
[DE/DE]; Bückmannmühle 46, 45326 Essen (DE).  
BRÜGGEMEIER, Ulf [DE/DE]; Leysiefen 20, 42799  
Leichlingen (DE). GEERTS, Andreas [DE/DE]; Schuck-  
ertstr. 29, 42113 Wuppertal (DE).

(74) Common Representative: BAYER HEALTHCARE  
AG; Law and Patents, Patents and Licensing, 51368  
Leverkusen (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,  
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,  
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,  
MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,  
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR,  
TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**

— as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii)) for the following designations AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH,  
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES,  
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,  
MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH,  
PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN,  
TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO  
patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG,  
ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU,  
TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE,  
DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT,  
RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

**Published:**

— with international search report  
— before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

(88) Date of publication of the international search report:  
5 August 2004

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: DIAGNOSTICS AND THERAPEUTICS FOR DISEASES ASSOCIATED WITH HUMAN PHOSPHODIESTERASE  
4D (PDE4D)

(57) Abstract: The invention provides a human PDE4D which is associated with the disorders of the peripheral and central nervous  
system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders. The invention also provides  
assays for the identification of compounds useful in the treatment or prevention of disorders of the peripheral and central nervous  
system, cardiovascular diseases, hematological diseases, respiratory diseases and urological disorders. The invention also features  
compounds which bind to and/or activate or inhibit the activity of PDE4D as well as pharmaceutical compositions comprising such  
compounds.

WO 2004/042389 A3



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/11877

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12Q1/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>PEREZ-TORRES S ET AL: "Cellular expression of cAMP phosphodiesterase PDE4A, PDE4B and PDE4D mRNAs in Alzheimer's disease brains" SOCIETY FOR NEUROSCIENCE ABSTRACTS, vol. 27, no. 1, 2001, page 1134, XP0002270742 &amp; 31ST ANNUAL MEETING OF THE SOCIETY FOR NEUROSCIENCE; SAN DIEGO, CALIFORNIA, USA; NOVEMBER 10-15, 2001 ISSN: 0190-5295 abstract</p> <p style="text-align: center;">----- -/-</p>	1-26

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

4 May 2004

Date of mailing of the international search report

26/05/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Thiele, U

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>KOSTIC M M ET AL: "Altered expression of PDE1 and PDE4 cyclic nucleotide phosphodiesterase isoforms in 7-oxo-prostacyclin-preconditioned rat heart."</p> <p>JOURNAL OF MOLECULAR AND CELLULAR CARDIOLOGY. NOV 1997, vol. 29, no. 11, November 1997 (1997-11), pages 3135-3146, XP002278927 ISSN: 0022-2828 abstract; paragraph bridging page 3141, r. h. col. and page 3142, r. h. col.</p>	1-26
X	<p>PEREZ-TORRES S ET AL: "Phosphodiesterase type 4 isozymes expression in human brain examined by in situ hybridization histochemistry and (3H)rolipram binding autoradiography comparison with monkey and rat brain"</p> <p>JOURNAL OF CHEMICAL NEUROANATOMY, CHICHESTER, GB, vol. 20, no. 3-4, December 2000 (2000-12), pages 349-374, XP002275597 ISSN: 0891-0618 abstract; table 1</p>	1-26
X	<p>MULLER T ET AL: "Subtypes of the type 4 cAMP phosphodiesterases: structure, regulation and selective inhibition"</p> <p>TRENDS IN PHARMACOLOGICAL SCIENCES, ELSEVIER TRENDS JOURNAL, CAMBRIDGE, GB, vol. 17, no. 8, 1 August 1996 (1996-08-01), pages 294-298, XP004034578 ISSN: 0165-6147 table 3</p>	1-26
X	<p>HANSEN G ET AL: "Absence of muscarinic cholinergic airway responses in mice deficient in the cyclic nucleotide phosphodiesterase PDE4D."</p> <p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA. 6 JUN 2000, vol. 97, no. 12, 6 June 2000 (2000-06-06), pages 6751-6756, XP002278928 ISSN: 0027-8424 abstract</p>	1-26

-/-

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/11877

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>LUGNIER C ET AL: "Characterization of cyclic nucleotide phosphodiesterase isoforms associated to isolated cardiac nuclei."  BIOCHIMICA ET BIOPHYSICA ACTA. 16 NOV 1999,  vol. 1472, no. 3,  16 November 1999 (1999-11-16), pages 431-446, XP002278929  ISSN: 0006-3002  abstract</p>	1-26
X	<p>GIEMBYCZ M: "PDE4D-deficient mice knock the breath out of asthma"  TRENDS IN PHARMACOLOGICAL SCIENCES, ELSEVIER TRENDS JOURNAL, CAMBRIDGE, GB, vol. 21, no. 8,  1 August 2000 (2000-08-01), pages 291-292, XP004215216  ISSN: 0165-6147  the whole document</p>	1-26
X	<p>LANDELLS L J ET AL: "IDENTIFICATION AND QUANTIFICATION OF PHOSPHODIESTERASE 4 SUBTYPES IN CD4 AND CD18 LYMPHOCYTES FROM HEALTHY AND ASTHMATIC SUBJECTS"  BRITISH JOURNAL OF PHARMACOLOGY, BASINGSTOKE, HANTS, GB,  vol. 133, no. 5, July 2001 (2001-07), pages 722-729, XP001037833  ISSN: 0007-1188  abstract; figure 1</p>	1-26
X	<p>WO 00/40714 A (ARROW AMY ; OLIGOS ETC INC (US); THOMPSON TERRY (US); DALE RODERIC M K) 13 July 2000 (2000-07-13)  pages 3, 17, 22 et seq. (Examples)</p>	1-26
A	<p>THOMAS N J ET AL: "Chronic type IV phosphodiesterase inhibition protects glomerular filtration rate and renal and mesenteric blood flow in a zymosan-induced model of multiple organ dysfunction syndrome treated with norepinephrine."  THE JOURNAL OF PHARMACOLOGY AND EXPERIMENTAL THERAPEUTICS. JAN 2001, vol. 296, no. 1, January 2001 (2001-01), pages 168-174, XP002278930  ISSN: 0022-3565  abstract</p>	
	-/--	

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/11877

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>TAKAHASHI KOKI ET AL: "Enhanced activities and gene expression of phosphodiesterase types 3 and 4 in pressure-induced congestive heart failure." HEART AND VESSELS. SEP 2002, vol. 16, no. 6, September 2002 (2002-09), pages 249-256, XP002278931 ISSN: 0910-8327 abstract; page 253 "PDE gene expression -----</p>	
A	<p>EP 1 040 829 A (PFIZER) 4 October 2000 (2000-10-04) claim 1 -----</p>	

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP 03/11877

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claim 26 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International Application No

PCT/EP 03/11877

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0040714	A	13-07-2000	AU 2480800 A	24-07-2000
			CA 2357950 A1	13-07-2000
			EP 1141278 A2	10-10-2001
			JP 2002534086 T	15-10-2002
			US 2003045490 A1	06-03-2003
			WO 0040714 A2	13-07-2000
<hr/>				
EP 1040829	A	04-10-2000	EP 1040829 A2	04-10-2000
			JP 2000198734 A	18-07-2000
<hr/>				